

Stochastic Lagrangian path for Leray solutions of 3D Navier-Stokes equations

Xicheng Zhang

Wuhan University

(A joint work with Guohuan Zhao)

Changchun • 2019.7.14

- Let \mathbf{u} be any Leray's solution of 3D-NSE

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi.$$

- Let \mathbf{u} be any Leray's solution of 3D-NSE

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi.$$

- We show the existence of weak solutions to the following stochastic Lagrangian particle equation

$$dX_{s,t} = \mathbf{u}(t, X_{s,t})dt + \sqrt{2\nu}dW_t, \quad X_{s,s} = x, \quad t \geq s.$$

- Let \mathbf{u} be any Leray's solution of 3D-NSE

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi.$$

- We show the existence of weak solutions to the following stochastic Lagrangian particle equation

$$dX_{s,t} = \mathbf{u}(t, X_{s,t})dt + \sqrt{2\nu}dW_t, \quad X_{s,s} = x, \quad t \geq s.$$

- $\mathbb{P} \circ X_{s,t}(x)^{-1} \in \mathbb{H}_q^{1,p}$ provided $p, q \in [1, 2)$ with $\frac{3}{p} + \frac{2}{q} > 4$.

- Let \mathbf{u} be any Leray's solution of 3D-NSE

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi.$$

- We show the existence of weak solutions to the following stochastic Lagrangian particle equation

$$dX_{s,t} = \mathbf{u}(t, X_{s,t})dt + \sqrt{2\nu}dW_t, \quad X_{s,s} = x, \quad t \geq s.$$

- $\mathbb{P} \circ X_{s,t}(x)^{-1} \in \mathbb{H}_q^{1,p}$ provided $p, q \in [1, 2)$ with $\frac{3}{p} + \frac{2}{q} > 4$.
- For Lebesgue almost all (s, x) , the solution $X_{s,\cdot}^n(x)$ associated with the mollifying velocity field \mathbf{u}_n weakly converges to $X_{s,\cdot}(x)$.

- 1 Introduction
- 2 Main result
- 3 Idea of Proof
- 4 Maximal principle by De-Giorgi's argument

- Let $d \geq 2$ and consider the following Navier-Stokes equation:

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi,$$

where $\mathbf{u} = (u_1, \dots, u_d)$ is the velocity field of the fluid, $\nu > 0$ is the viscosity constant, and p stands for the pressure.

- Let $d \geq 2$ and consider the following Navier-Stokes equation:

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi,$$

where $\mathbf{u} = (u_1, \dots, u_d)$ is the velocity field of the fluid, $\nu > 0$ is the viscosity constant, and p stands for the pressure.

- For any divergence free vector field $\varphi \in L^2(\mathbb{R}^d; \mathbb{R}^d)$, there exists a divergence free Leray weak solution to 3D-NSEs with

$$\|\mathbf{u}\|_{L^\infty([0, T]; L^2(\mathbb{R}^d))} + \|\nabla \mathbf{u}\|_{L^2([0, T]; L^2(\mathbb{R}^d))} < \infty, \quad \forall T > 0.$$

- Let $d \geq 2$ and consider the following Navier-Stokes equation:

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi,$$

where $\mathbf{u} = (u_1, \dots, u_d)$ is the velocity field of the fluid, $\nu > 0$ is the viscosity constant, and p stands for the pressure.

- For any divergence free vector field $\varphi \in L^2(\mathbb{R}^d; \mathbb{R}^d)$, there exists a divergence free Leray weak solution to 3D-NSEs with

$$\|\mathbf{u}\|_{L^\infty([0, T]; L^2(\mathbb{R}^d))} + \|\nabla \mathbf{u}\|_{L^2([0, T]; L^2(\mathbb{R}^d))} < \infty, \quad \forall T > 0.$$

- Buckmaster and Vicol (2018, AOM)** showed that there are **infinitely many** weak solutions $\mathbf{u} \in C(\mathbb{R}_+; L^2(\mathbb{T}^3))$ for 3D-NSEs on the torus.

- Let $d \geq 2$ and consider the following Navier-Stokes equation:

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p, \quad \operatorname{div} \mathbf{u} \equiv 0, \quad \mathbf{u}_0 = \varphi,$$

where $\mathbf{u} = (u_1, \dots, u_d)$ is the velocity field of the fluid, $\nu > 0$ is the viscosity constant, and p stands for the pressure.

- For any divergence free vector field $\varphi \in L^2(\mathbb{R}^d; \mathbb{R}^d)$, there exists a divergence free Leray weak solution to 3D-NSEs with

$$\|\mathbf{u}\|_{L^\infty([0, T]; L^2(\mathbb{R}^d))} + \|\nabla \mathbf{u}\|_{L^2([0, T]; L^2(\mathbb{R}^d))} < \infty, \quad \forall T > 0.$$

- Buckmaster and Vicol (2018, AOM)** showed that there are **infinitely many** weak solutions $\mathbf{u} \in C(\mathbb{R}_+; L^2(\mathbb{T}^3))$ for 3D-NSEs on the torus.
- Existence and smoothness of Leray solutions are **open problems!**

- Question: For any Leray solution \mathbf{u} , is it possible to construct the stochastic Lagrangian particle trajectory associated with \mathbf{u} ?

- Question: For any Leray solution \mathbf{u} , is it possible to construct the **stochastic Lagrangian particle trajectory** associated with \mathbf{u} ?
- More precisely, for each starting point x , is there a unique solution to the following SDE?

$$dX_t = \mathbf{u}(t, X_t)dt + \sqrt{2\nu}dW_t, \quad X_0 = x, \quad (1.1)$$

where W is a d -dimensional standard Brownian motion on some probability space $(\Omega, \mathcal{F}, \mathbf{P})$.

- If \mathbf{u} is smooth in x , then by [Constantin and Iyer's representation \(2008, CPAM\)](#), \mathbf{u} can be reconstructed from $X_t(x)$ as follows:

$$\mathbf{u}(t, x) = \mathcal{P}\mathbb{E}(\nabla^t X_t^{-1}(x) \cdot \varphi(X_t^{-1}(x))),$$

where \mathcal{P} is the Leray projection and $X_t^{-1}(x)$ is the inverse of stochastic flow $x \mapsto X_t(x)$, and ∇^t is the transpose of a Jacobian matrix.

- Krylov and Röckner (2005, PTRF) showed the existence-uniqueness of strong solutions to SDE (1.1) under the following assumption

$$\mathbf{u} \in \cap_{T>0} L^q([0, T]; L^p(\mathbb{R}^d)), \quad p, q \geq 2, \quad \frac{d}{p} + \frac{2}{q} < 1. \quad (1.2)$$

- Krylov and Röckner (2005, PTRF) showed the existence-uniqueness of strong solutions to SDE (1.1) under the following assumption

$$\mathbf{u} \in \cap_{T>0} L^q([0, T]; L^p(\mathbb{R}^d)), \quad p, q \geq 2, \quad \frac{d}{p} + \frac{2}{q} < 1. \quad (1.2)$$

- The unique solution $X_t(x)$ is weakly differentiable in x and satisfies (see Fedrizzi-Flandoli (2013), Z. (2013), (2016)):

$$\sup_{x \in \mathbb{R}^d} \mathbf{E} \left(\sup_{t \in [0, T]} |\nabla X_t(x)|^p \right) < \infty, \quad \forall p \geq 1, \quad T > 0.$$

- Leray's solution does not satisfy

$$\mathbf{u} \in \cap_{T>0} L^q([0, T]; L^p(\mathbb{R}^d)), \quad p, q \geq 2, \quad \frac{d}{p} + \frac{2}{q} < 1, \quad (1.3)$$

but only satisfies

$$\mathbf{u} \in \cap_{T>0} L^q([0, T]; L^p(\mathbb{R}^d)), \quad p, q \geq 2, \quad \frac{d}{p} + \frac{2}{q} > \frac{d}{2}. \quad (1.4)$$

- Leray's solution does not satisfy

$$\mathbf{u} \in \cap_{T>0} L^q([0, T]; L^p(\mathbb{R}^d)), \quad p, q \geq 2, \quad \frac{d}{p} + \frac{2}{q} < 1, \quad (1.3)$$

but only satisfies

$$\mathbf{u} \in \cap_{T>0} L^q([0, T]; L^p(\mathbb{R}^d)), \quad p, q \geq 2, \quad \frac{d}{p} + \frac{2}{q} > \frac{d}{2}. \quad (1.4)$$

- **Deterministic** Lagrangian particle trajectories associated with \mathbf{u} have been studied very well (see [Robinson, Rodrigo and Sadowski's](#) book).

- Consider the following stochastic differential equation in \mathbb{R}^d :

$$dX_{s,t} = b(t, X_{s,t})dt + \sqrt{2}dW_t, \quad t > s, \quad X_{s,s} = x, \quad (1.5)$$

where $b(t, x) : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a measurable vector field.

- Consider the following stochastic differential equation in \mathbb{R}^d :

$$dX_{s,t} = b(t, X_{s,t})dt + \sqrt{2}dW_t, \quad t > s, \quad X_{s,s} = x, \quad (1.5)$$

where $b(t, x) : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a measurable vector field.

- The generator associated with the above SDE is given by

$$\mathcal{L}_t^b := \Delta + b(t, \cdot) \cdot \nabla.$$

- Consider the following stochastic differential equation in \mathbb{R}^d :

$$dX_{s,t} = b(t, X_{s,t})dt + \sqrt{2}dW_t, \quad t > s, \quad X_{s,s} = x, \quad (1.5)$$

where $b(t, x) : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a measurable vector field.

- The generator associated with the above SDE is given by

$$\mathcal{L}_t^b := \Delta + b(t, \cdot) \cdot \nabla.$$

- Under the following assumption

$$b \in \cap_{T>0} L^q([0, T]; L^p(\mathbb{R}^d)) =: L_{loc}^q(L^p), \quad p, q \geq 2, \quad \frac{d}{p} + \frac{2}{q} < 2,$$

is there a semimartingale solution of SDE (1.5)? That is,

$$X_{s,t} = x + \int_s^t b(r, X_{s,r})dr + \sqrt{2}(W_t - W_s), \quad \forall t \geq s ?? \quad (1.6)$$

- Krylov-Röckner (2005, PTRF): Strong well-posedness ($\frac{d}{p} + \frac{2}{q} < 1$).

- Krylov-Röckner (2005, PTRF): Strong well-posedness ($\frac{d}{p} + \frac{2}{q} < 1$).
- Bass-Chen (2006, AOP): Weak well-posedness in the class of semimartingales when b belongs to some generalized Kato's class \mathbf{K}_{d-1} .

- Krylov-Röckner (2005, PTRF): Strong well-posedness ($\frac{d}{p} + \frac{2}{q} < 1$).
- Bass-Chen (2006, AOP): Weak well-posedness in the class of semimartingales when b belongs to some generalized Kato's class \mathbf{K}_{d-1} .
- Flandoli, Issoglio and Russo (2014) Well-posedness in the class of “virtual” solutions when $b \in H^{-\alpha,p}$ with $\alpha \in (0, \frac{1}{2})$ and $p \in (\frac{d}{1-\alpha}, \frac{d}{\alpha})$.

- Krylov-Röckner (2005, PTRF): Strong well-posedness ($\frac{d}{p} + \frac{2}{q} < 1$).
- Bass-Chen (2006, AOP): Weak well-posedness in the class of semimartingales when b belongs to some generalized Kato's class \mathbf{K}_{d-1} .
- Flandoli, Issoglio and Russo (2014) Well-posedness in the class of “virtual” solutions when $b \in H^{-\alpha,p}$ with $\alpha \in (0, \frac{1}{2})$ and $p \in (\frac{d}{1-\alpha}, \frac{d}{\alpha})$.
- Z.-Zhao (2017): Weak well-posedness in the class of Dirichlet processes when $b \in H^{-\alpha,p}$ with $\alpha \in (0, \frac{1}{2}]$ and $p \in (\frac{d}{1-\alpha}, \infty)$.
-

- Critical case ($\frac{d}{p} + \frac{2}{q} = 1$ with $p, q \in (2, \infty)$):

Beck, Flandoli, Gubinelli and Maurelli (2014) showed the existence of strong solutions for almost all starting point $x \in \mathbb{R}^d$.

- Critical case ($\frac{d}{p} + \frac{2}{q} = 1$ with $p, q \in (2, \infty)$):

Beck, Flandoli, Gubinelli and Maurelli (2014) showed the existence of strong solutions for almost all starting point $x \in \mathbb{R}^d$.

- Kinzebulatov and Semenov (2017) showed the existence of weak solutions for each starting point $x \in \mathbb{R}^d$ when $b \in L^d(\mathbb{R}^d)$ is *time-independent*, but the uniqueness is left open.

A counter-example

- Consider the following concrete SDE:

$$X_t = -c \int_0^t X_s |X_s|^{-2} ds + W_t, \quad c \in \mathbb{R}. \quad (1.7)$$

A counter-example

- Consider the following concrete SDE:

$$X_t = -c \int_0^t X_s |X_s|^{-2} ds + W_t, \quad c \in \mathbb{R}. \quad (1.7)$$

- If $c \geq d$, [Kinzebulatov and Semenov](#) showed that the above SDE does not allow a solution. If $c < c_d$, where $c_d \in (0, d)$ is some constant only depending on d , they proved that there exists a weak solution to the above SDE by utilizing the analytic construction of the semigroup $e^{-t(\Delta + b \cdot \nabla)}$.

A counter-example

- Consider the following concrete SDE:

$$X_t = -c \int_0^t X_s |X_s|^{-2} ds + W_t, \quad c \in \mathbb{R}. \quad (1.7)$$

- If $c \geq d$, [Kinzebulatov and Semenov](#) showed that the above SDE does not allow a solution. If $c < c_d$, where $c_d \in (0, d)$ is some constant only depending on d , they proved that there exists a weak solution to the above SDE by utilizing the analytic construction of the semigroup $e^{-t(\Delta + b \cdot \nabla)}$.
- By direct calculations, for $b(x) := -cx|x|^{-2}$ and $d \geq 3$, we have

$$\operatorname{div} b(x) = -c(d-2)|x|^{-2} \notin L_{loc}^{d/2}.$$

A counter-example

- Consider the following concrete SDE:

$$X_t = -c \int_0^t X_s |X_s|^{-2} ds + W_t, \quad c \in \mathbb{R}. \quad (1.7)$$

- If $c \geq d$, [Kinzebulatov and Semenov](#) showed that the above SDE does not allow a solution. If $c < c_d$, where $c_d \in (0, d)$ is some constant only depending on d , they proved that there exists a weak solution to the above SDE by utilizing the analytic construction of the semigroup $e^{-t(\Delta + b \cdot \nabla)}$.
- By direct calculations, for $b(x) := -cx|x|^{-2}$ and $d \geq 3$, we have

$$\operatorname{div} b(x) = -c(d-2)|x|^{-2} \notin L_{loc}^{d/2}.$$

- Intuitively, if $c \geq d$, then the centripetal force is so strong such that the particle can not escape from the origin immediately so that even though a random perturbation is added, there is no solution for SDE (1.7).

Main result

- Let $C_c^\infty(\mathbb{R}^{d+1})$ be the space of all smooth functions with compact supports and \mathcal{D}' the dual space of $C_c^\infty(\mathbb{R}^{d+1})$, which is also called distribution space. The duality between \mathcal{D}' and $C_c^\infty(\mathbb{R}^{d+1})$ is denoted by $\langle\langle \cdot, \cdot \rangle\rangle$. In particular, if $f(t, x)$ and $g(t, x)$ are two real functions in $\mathbb{R} \times \mathbb{R}^d$, then

$$\langle\langle f, g \rangle\rangle = \int_{\mathbb{R}} \langle f(t), g(t) \rangle dt \quad \text{with} \quad \langle f(t), g(t) \rangle := \int_{\mathbb{R}^d} f(t, x)g(t, x)dx.$$

Main result

- Let $C_c^\infty(\mathbb{R}^{d+1})$ be the space of all smooth functions with compact supports and \mathcal{D}' the dual space of $C_c^\infty(\mathbb{R}^{d+1})$, which is also called distribution space. The duality between \mathcal{D}' and $C_c^\infty(\mathbb{R}^{d+1})$ is denoted by $\langle\langle \cdot, \cdot \rangle\rangle$. In particular, if $f(t, x)$ and $g(t, x)$ are two real functions in $\mathbb{R} \times \mathbb{R}^d$, then

$$\langle\langle f, g \rangle\rangle = \int_{\mathbb{R}} \langle f(t), g(t) \rangle dt \quad \text{with} \quad \langle f(t), g(t) \rangle := \int_{\mathbb{R}^d} f(t, x)g(t, x)dx.$$

- For two distributions $f, g \in \mathcal{D}'$, one says that $f \leq g$ if for any non-negative $\varphi \in C_c^\infty(\mathbb{R}^{d+1})$,

$$\langle\langle f, \varphi \rangle\rangle \leq \langle\langle g, \varphi \rangle\rangle.$$

- For $\alpha \in \mathbb{R}$ and $p \in [1, \infty]$, let $H^{\alpha,p}$ be the usual Bessel potential space with norm:

$$\|f\|_{\alpha,p} := \|(\mathbb{I} - \Delta)^{\alpha/2} f\|_p = \left(\int_{\mathbb{R}^d} |(\mathbb{I} - \Delta)^{\alpha/2} f(x)|^p dx \right)^{1/p}.$$

- For $\alpha \in \mathbb{R}$ and $p \in [1, \infty]$, let $H^{\alpha,p}$ be the usual Bessel potential space with norm:

$$\|f\|_{\alpha,p} := \|(\mathbb{I} - \Delta)^{\alpha/2} f\|_p = \left(\int_{\mathbb{R}^d} |(\mathbb{I} - \Delta)^{\alpha/2} f(x)|^p dx \right)^{1/p}.$$

- For $\alpha \in \mathbb{R}$ and $p, q \in [1, \infty]$, let $\mathbb{H}_q^{\alpha,p} := L^q(\mathbb{R}; H^{\alpha,p})$ be the space of spatial-time functions with norm

$$\|f\|_{\alpha,p;q} := \left(\int_{\mathbb{R}} \|f(t, \cdot)\|_{\alpha,p}^q dt \right)^{1/q}.$$

- For $r > 0$, we define

$$B_r := \{x \in \mathbb{R}^d : |x| < r\}, \quad Q_r := (-r^2, r^2) \times B_r.$$

- For $r > 0$, we define

$$B_r := \{x \in \mathbb{R}^d : |x| < r\}, \quad Q_r := (-r^2, r^2) \times B_r.$$

- Fix $\chi \in C^\infty(\mathbb{R}^{d+1}; [0, 1])$ with $\chi|_{Q_1} = 1$ and $\chi|_{Q_2^c} = 0$. For $r > 0$ and $(s, z) \in \mathbb{R}^{d+1}$, define

$$\chi_r(t, x) := \chi(r^{-2}t, r^{-1}x), \quad \chi_r^{s,z}(t, x) := \chi_r(t - s, x - z).$$

- For $r > 0$, we define

$$B_r := \{x \in \mathbb{R}^d : |x| < r\}, \quad Q_r := (-r^2, r^2) \times B_r.$$

- Fix $\chi \in C^\infty(\mathbb{R}^{d+1}; [0, 1])$ with $\chi|_{Q_1} = 1$ and $\chi|_{Q_2^c} = 0$. For $r > 0$ and $(s, z) \in \mathbb{R}^{d+1}$, define

$$\chi_r(t, x) := \chi(r^{-2}t, r^{-1}x), \quad \chi_r^{s,z}(t, x) := \chi_r(t - s, x - z).$$

- Fix $r > 0$. Let $\widetilde{\mathbb{H}}_q^{\alpha,p}$ be the Banach space of all functions $f \in \mathbb{H}_{q,loc}^{\alpha,p}$ with

$$\|f\|_{\alpha,p;q} := \sup_{s,z} \|f \chi_r^{s,z}\|_{\alpha,p;q} < \infty.$$

- For $r > 0$, we define

$$B_r := \{x \in \mathbb{R}^d : |x| < r\}, \quad Q_r := (-r^2, r^2) \times B_r.$$

- Fix $\chi \in C^\infty(\mathbb{R}^{d+1}; [0, 1])$ with $\chi|_{Q_1} = 1$ and $\chi|_{Q_2^c} = 0$. For $r > 0$ and $(s, z) \in \mathbb{R}^{d+1}$, define

$$\chi_r(t, x) := \chi(r^{-2}t, r^{-1}x), \quad \chi_r^{s,z}(t, x) := \chi_r(t - s, x - z).$$

- Fix $r > 0$. Let $\tilde{\mathbb{H}}_q^{\alpha,p}$ be the Banach space of all functions $f \in \mathbb{H}_{q,loc}^{\alpha,p}$ with

$$\|f\|_{\alpha,p;q} := \sup_{s,z} \|f \chi_r^{s,z}\|_{\alpha,p;q} < \infty.$$

- For $p' > p, q' > q$, we have $\tilde{\mathbb{H}}_{q'}^{\alpha,p'} \subset \tilde{\mathbb{H}}_q^{\alpha,p}$.

- Let \mathbb{C} be the space of all continuous functions from \mathbb{R}_+ to \mathbb{R}^d , which is endowed with the usual Borel σ -field $\mathcal{B}(\mathbb{C})$.

- Let \mathbb{C} be the space of all continuous functions from \mathbb{R}_+ to \mathbb{R}^d , which is endowed with the usual Borel σ -field $\mathcal{B}(\mathbb{C})$.
- All the probability measures over $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ is denoted by $\mathcal{P}(\mathbb{C})$.

- Let \mathbb{C} be the space of all continuous functions from \mathbb{R}_+ to \mathbb{R}^d , which is endowed with the usual Borel σ -field $\mathcal{B}(\mathbb{C})$.
- All the probability measures over $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ is denoted by $\mathcal{P}(\mathbb{C})$.
- Let ω_t be the canonical process over \mathbb{C} . For $t \geq 0$, let $\mathcal{B}_t := \mathcal{B}_t(\mathbb{C})$ be the natural filtration generated by $\{\omega_s : s \leq t\}$.

Definition 1

For given $(s, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, we call a probability measure $\mathbb{P}_{s,x} \in \mathcal{P}(\mathbb{C})$ a martingale solution of SDE (1.5) with starting point (s, x) if

(i) $\mathbb{P}_{s,x}(\omega_t = x, t \leq s) = 1$, and for each $t > s$,

$$\mathbb{E}^{\mathbb{P}_{s,x}} \left(\int_s^t |b(r, \omega_r)| dr \right) < \infty.$$

(ii) For all $f \in C_c^2(\mathbb{R}^d)$, M_t^f is a \mathcal{B}_t -martingale under $\mathbb{P}_{s,x}$, where

$$M_t^f(\omega) := f(\omega_t) - f(x) - \int_s^t \mathcal{L}_r^b f(\omega_r) dr, \quad t \geq s.$$

All the martingale solution $\mathbb{P}_{s,x}$ with starting point (s, x) and drift b is denoted by $\mathcal{M}_{s,x}^b$.

- Let $\mathbb{P}_{s,x} \in \mathcal{M}_{s,x}^b$. By Lévy's characterization for Brownian motion, one sees that

$$W_t := \frac{\sqrt{2}}{2} \left(\omega_t - \omega_s - \int_s^t b(r, \omega_r) dr \right), \quad t \geq s,$$

is a d -dimensional standard Brownian motion under $\mathbb{P}_{s,x}$, so that

$$\omega_t = x + \int_s^t b(r, \omega_r) dr + \sqrt{2} W_t, \quad t \geq s.$$

In other words, $(\mathbb{C}, \mathcal{B}(\mathbb{C}), \mathbb{P}_{s,x}, \omega_t, W_t)$ is a weak solution of SDE (1.5).

Theorem 2

Suppose that for some $p_i, q_i \in [2, \infty)$ with $\frac{d}{p_i} + \frac{2}{q_i} < 2$, $i = 1, 2$,

$$\| \|b\| \|_{0,p_1;q_1} + \| \|(\operatorname{div} b)^-\| \|_{0,p_2;q_2} < \infty. \quad (2.1)$$

For each $(s, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, there exists at least one martingale solution $\mathbb{P}_{s,x} \in \mathcal{M}_{s,x}^b$, which satisfies the following Krylov's type estimate: for any $\alpha \in [0, 1]$ and $p, q \in (1, \infty)$ with $\frac{d}{p} + \frac{2}{q} < 2 - \alpha$, there exist $\theta = \theta(\alpha, p, q) > 0$ and a constant $C > 0$ such that for all $s \leq t_0 < t_1 < \infty$ with $t_1 - t_0 \leq 1$ and $f \in C_c^\infty(\mathbb{R}^{d+1})$,

$$\mathbb{E}^{\mathbb{P}_{s,x}} \left(\int_{t_0}^{t_1} f(t, \omega_t) dt \middle| \mathcal{B}_{t_0} \right) \leq C(t_1 - t_0)^\theta \| \|f\| \|_{-\alpha,p;q}. \quad (2.2)$$

Moreover, we have the following conclusions

- (i) (Weak uniqueness) For any mollifying approximation b_n of b , there is a Lebesgue-null set $\mathcal{N} \subset \mathbb{R}_+ \times \mathbb{R}^d$ such that for all $(s, x) \in \mathcal{N}^c$,

$$\mathbb{P}_{s,x}^n \text{ weakly converges to } \mathbb{P}_{s,x} \in \mathcal{M}_{s,x}^b, \text{ where } \mathbb{P}_{s,x}^n \in \mathcal{M}_{s,x}^{b_n}.$$

- (ii) (Almost surely Markov property) For each $(s, x) \in \mathcal{N}^c$, there is a Lebesgue null set $I_{s,x} \subset [s, \infty)$ such that for all $t_0 \in (s, \infty) \setminus I_{s,x}$, any $t_1 > t_0$ and $f \in C_c(\mathbb{R}^d)$,

$$\mathbb{E}^{\mathbb{P}_{s,x}}(f(\omega_{t_1}) | \mathcal{B}_{t_0}) = \mathbb{E}^{\mathbb{P}_{t_0, \omega_{t_0}}}(f(\omega_{t_1})), \quad \mathbb{P}_{s,x} - \text{a.s.}$$

- (iii) (L^p -semigroup) Let $\mathcal{T}_{s,t}f(x) := \mathbb{E}^{\mathbb{P}_{s,x}}f(\omega_t)$. For any $p \geq 1$ and $T > 0$, there is a constant $C > 0$ such that for Lebesgue almost all $0 \leq s < t \leq T$ and $f \in L^p(\mathbb{R}^d)$,

$$\|\mathcal{T}_{s,t}f\|_p \leq C\|f\|_p. \quad (2.3)$$

Application to stochastic Lagrangian particle path of 3D-NSEs

- If $(\operatorname{div} b)^- \equiv 0$, then $\|\mathcal{T}_{s,t}f\|_1 \leq \|f\|_1$ in (2.3). If $\operatorname{div} b \equiv 0$, then for any nonnegative $f \in L^1(\mathbb{R}^d)$, $\|\mathcal{T}_{s,t}f\|_1 = \|f\|_1$. By (1.4), we can apply the above theorem to the Leray solution of 3D-NSEs.

Application to stochastic Lagrangian particle path of 3D-NSEs

- If $(\operatorname{div} b)^- \equiv 0$, then $\|\mathcal{T}_{s,t} f\|_1 \leq \|f\|_1$ in (2.3). If $\operatorname{div} b \equiv 0$, then for any nonnegative $f \in L^1(\mathbb{R}^d)$, $\|\mathcal{T}_{s,t} f\|_1 = \|f\|_1$. By (1.4), we can apply the above theorem to the Leray solution of 3D-NSEs.
- Under the assumptions

$$\nabla b \in \mathbb{L}_{loc}^1, \quad (\operatorname{div} b)^-, b/(1 + |x|) \in \mathbb{L}^\infty,$$

the existence and uniqueness of almost everywhere stochastic flows are obtained in the framework of DiPerna-Lions' theory has been obtained in Zhang (2010). However, the existence of a solution is only shown for Lebesgue almost all starting point $x \in \mathbb{R}^d$.

Example

Let $d \geq 3$ and $\alpha < 3$. Define

$$b(x) := \sum_{z \in \mathbb{Z}^d} \gamma_z \frac{x - z}{|x - z|^\alpha} \phi(|x - z|),$$

where for some $M > 0$, $\gamma_z \in (0, M)$ is a constant and $\phi \in C_c^\infty(\mathbb{R}_+; [0, 1])$ with $\phi(r) = 1$ for $r \in [0, 1]$ and $\phi(r) = 0$ for $r > 2$. It is easy to see that (2.1) holds.

Idea of Proof

- We assume that for some $p_i, q_i \in [2, \infty)$ with $\frac{d}{p_i} + \frac{2}{q_i} < 2$, $i = 1, 2$,

$$\kappa := \|b\|_{p_1; q_1} + \|(\operatorname{div} b)^-\|_{p_2; q_2} < \infty.$$

Idea of Proof

- We assume that for some $p_i, q_i \in [2, \infty)$ with $\frac{d}{p_i} + \frac{2}{q_i} < 2, i = 1, 2,$

$$\kappa := \|\|b\|\|_{p_1; q_1} + \|\|(\operatorname{div} b)^-\|\|_{p_2; q_2} < \infty.$$

- Let $b_n(t, x) = b(t, \cdot) * \rho_n(x)$ be the mollifying approximation of $b(t, \cdot)$. It is easy to check that

$$\sup_n (\|\|b_n\|\|_{p_1; q_1} + \|\|(\operatorname{div} b_n)^-\|\|_{p_2; q_2}) \leq C\kappa,$$

and

$$b_n \in L_{loc}^{q_1}(\mathbb{R}_+; C_b^\infty(\mathbb{R}^d)).$$

Idea of Proof

- We assume that for some $p_i, q_i \in [2, \infty)$ with $\frac{d}{p_i} + \frac{2}{q_i} < 2, i = 1, 2,$

$$\kappa := |||b|||_{p_1; q_1} + |||(\operatorname{div} b)^-|||_{p_2; q_2} < \infty.$$

- Let $b_n(t, x) = b(t, \cdot) * \rho_n(x)$ be the mollifying approximation of $b(t, \cdot)$. It is easy to check that

$$\sup_n (|||b_n|||_{p_1; q_1} + |||(\operatorname{div} b_n)^-|||_{p_2; q_2}) \leq C\kappa,$$

and

$$b_n \in L_{loc}^{q_1}(\mathbb{R}_+; C_b^\infty(\mathbb{R}^d)).$$

- For $(s, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, consider the following SDE:

$$dX_{s,t}^n = b_n(t, X_{s,t}^n)dt + \sqrt{2}dW_t, \quad X_{s,s}^n = x, \quad t \geq s,$$

where W is a d -dimensional standard Brownian motion on some complete filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$.

- We want to show that the law $\mathbb{P}_{s,x}^n$ of $X_{s,\cdot}^n(x)$ is tight in \mathbb{C} .

- We want to show that the law $\mathbb{P}_{s,x}^n$ of $X_{s,\cdot}^n(x)$ is tight in \mathbb{C} .
- Notice that for any stopping time τ ,

$$X_{s,\tau+\delta}^n(x) - X_{s,\tau}^n(x) = \int_{\tau}^{\tau+\delta} b_n(t, X_{s,t}^n(x)) dt + \sqrt{2}(W_{\tau+\delta} - W_{\tau}), \quad \delta > 0.$$

- We want to show that the law $\mathbb{P}_{s,x}^n$ of $X_{s,\cdot}^n(x)$ is tight in \mathbb{C} .
- Notice that for any stopping time τ ,

$$X_{s,\tau+\delta}^n(x) - X_{s,\tau}^n(x) = \int_{\tau}^{\tau+\delta} b_n(t, X_{s,t}^n(x)) dt + \sqrt{2}(W_{\tau+\delta} - W_{\tau}), \quad \delta > 0.$$

- If we can show that for some $\alpha > 0$,

$$\lim_{\delta \rightarrow 0} \sup_n \sup_{\tau} \mathbb{E} |X_{s,\tau+\delta}^n(x) - X_{s,\tau}^n(x)|^{\alpha} = 0,$$

then the tightness follows.

- We want to show that the law $\mathbb{P}_{s,x}^n$ of $X_{s,\cdot}^n(x)$ is tight in \mathbb{C} .
- Notice that for any stopping time τ ,

$$X_{s,\tau+\delta}^n(x) - X_{s,\tau}^n(x) = \int_{\tau}^{\tau+\delta} b_n(t, X_{s,t}^n(x)) dt + \sqrt{2}(W_{\tau+\delta} - W_{\tau}), \quad \delta > 0.$$

- If we can show that for some $\alpha > 0$,

$$\lim_{\delta \rightarrow 0} \sup_n \sup_{\tau} \mathbb{E} |X_{s,\tau+\delta}^n(x) - X_{s,\tau}^n(x)|^{\alpha} = 0,$$

then the tightness follows.

- By the strong Markov property, it suffices to show

$$\sup_n \sup_{s,x} \mathbb{E} \int_0^{\delta} b_n(s+t, X_{s,t}^n(x)) dt \leq c_{\delta} \rightarrow 0.$$

Lemma 3

For any $\alpha \in [0, 1]$ and $p, q \in (1, \infty)$ with $\frac{d}{p} + \frac{2}{q} < 2 - \alpha$, there are constants $\theta = \theta(\alpha, p, q) > 0$ and $C > 0$ depending on $\kappa, d, \alpha, p, q, p_i, q_i$ such that for any $f \in C_c^\infty(\mathbb{R}^{d+1})$ and $0 \leq s \leq t_0 < t_1 < \infty$ with $t_1 - t_0 \leq 1$,

$$\sup_n \sup_{x \in \mathbb{R}^d} \mathbf{E} \left(\int_{t_0}^{t_1} f(t, X_{s,t}^n(x)) dt \middle| \mathcal{F}_{t_0} \right) \leq C(t_1 - t_0)^\theta \|f\|_{-\alpha, p; q}.$$

- Fix $0 \leq s \leq t_0 < t_1 < \infty$ with $t_1 - t_0 \leq 1$ and $f \in C_c^\infty(\mathbb{R}^{d+1})$. Let u_n be the smooth solution of the following backward PDE:

$$\partial_t u_n + \Delta u_n + b_n \cdot \nabla u_n + f = 0, \quad u_n(t_1, \cdot) = 0.$$

- Fix $0 \leq s \leq t_0 < t_1 < \infty$ with $t_1 - t_0 \leq 1$ and $f \in C_c^\infty(\mathbb{R}^{d+1})$. Let u_n be the smooth solution of the following backward PDE:

$$\partial_t u_n + \Delta u_n + b_n \cdot \nabla u_n + f = 0, \quad u_n(t_1, \cdot) = 0.$$

- By Itô's formula we have

$$u_n(t_1, X_{s,t_1}^n) = u_n(t_0, X_{s,t_0}^n) - \int_{t_0}^{t_1} f(t, X_{s,t}^n) dt + \sqrt{2} \int_{t_0}^{t_1} \nabla u_n(t, X_{s,t}^n) dW_t.$$

- Fix $0 \leq s \leq t_0 < t_1 < \infty$ with $t_1 - t_0 \leq 1$ and $f \in C_C^\infty(\mathbb{R}^{d+1})$. Let u_n be the smooth solution of the following backward PDE:

$$\partial_t u_n + \Delta u_n + b_n \cdot \nabla u_n + f = 0, \quad u_n(t_1, \cdot) = 0.$$

- By Itô's formula we have

$$u_n(t_1, X_{s,t_1}^n) = u_n(t_0, X_{s,t_0}^n) - \int_{t_0}^{t_1} f(t, X_{s,t}^n) dt + \sqrt{2} \int_{t_0}^{t_1} \nabla u_n(t, X_{s,t}^n) dW_t.$$

- By taking conditional expectation with respect to \mathcal{F}_{t_0} , we obtain

$$\mathbf{E} \left(\int_{t_0}^{t_1} f(t, X_{s,t}^n) dt \middle| \mathcal{F}_{t_0} \right) = \mathbf{E} \left(u_n(t_0, X_{s,t_0}^n) \middle| \mathcal{F}_{t_0} \right) \leq \|u_n(t_0)\|_\infty.$$

- Fix $0 \leq s \leq t_0 < t_1 < \infty$ with $t_1 - t_0 \leq 1$ and $f \in C_c^\infty(\mathbb{R}^{d+1})$. Let u_n be the smooth solution of the following backward PDE:

$$\partial_t u_n + \Delta u_n + b_n \cdot \nabla u_n + f = 0, \quad u_n(t_1, \cdot) = 0.$$

- By Itô's formula we have

$$u_n(t_1, X_{s,t_1}^n) = u_n(t_0, X_{s,t_0}^n) - \int_{t_0}^{t_1} f(t, X_{s,t}^n) dt + \sqrt{2} \int_{t_0}^{t_1} \nabla u_n(t, X_{s,t}^n) dW_t.$$

- By taking conditional expectation with respect to \mathcal{F}_{t_0} , we obtain

$$\mathbf{E} \left(\int_{t_0}^{t_1} f(t, X_{s,t}^n) dt \middle| \mathcal{F}_{t_0} \right) = \mathbf{E} \left(u_n(t_0, X_{s,t_0}^n) \middle| \mathcal{F}_{t_0} \right) \leq \|u_n(t_0)\|_\infty.$$

- The key point is to show that

$$\|u_n(t_0)\|_\infty \leq C \|f\|_{[-\alpha, p; q']} \leq C (t_1 - t_0)^{1 - \frac{q'}{q}} \|f\|_{[-\alpha, p; q]},$$

where $q' < q$ so that $\frac{d}{p} + \frac{2}{q'} < 2 - \alpha$.

Maximal principle by De-Giorgi's argument

- Let $\mathcal{V} := \mathbb{L}_{\infty}^2 \cap \mathbb{H}_2^{1,2}$, $\mathcal{V}_{loc} := \mathbb{L}_{\infty,loc}^2 \cap \mathbb{H}_{2,loc}^{1,2}$.

Maximal principle by De-Giorgi's argument

- Let $\mathcal{V} := \mathbb{L}_{\infty}^2 \cap \mathbb{H}_2^{1,2}$, $\mathcal{V}_{loc} := \mathbb{L}_{\infty,loc}^2 \cap \mathbb{H}_{2,loc}^{1,2}$.
- We assume

$$b \in \mathbb{L}_{2,loc}^2, \quad f \in \mathcal{D}',$$

Maximal principle by De-Giorgi's argument

- Let $\mathcal{V} := \mathbb{L}_{\infty}^2 \cap \mathbb{H}_2^{1,2}$, $\mathcal{V}_{loc} := \mathbb{L}_{\infty,loc}^2 \cap \mathbb{H}_{2,loc}^{1,2}$.
- We assume

$$\mathbf{b} \in \mathbb{L}_{2,loc}^2, \quad \mathbf{f} \in \mathcal{D}',$$

- Consider the following PDE in \mathbb{R}^{d+1} :

$$\partial_t u = \Delta u + \mathbf{b} \cdot \nabla u + f. \quad (4.1)$$

Definition 4

A function $u \in \mathcal{V}_{loc} \cap \mathbb{L}_{loc}^{\infty}$ is called a weak solution of PDE (4.1) if for any nonnegative smooth function $\varphi \in C_c^{\infty}(\mathbb{R}^{d+1})$ and almost all $t \in \mathbb{R}$,

$$\langle \partial_t u, \varphi \rangle = -\langle \nabla u, \nabla \varphi \rangle + \langle \mathbf{b} \cdot \nabla u, \varphi \rangle + \langle \mathbf{f}, \varphi \rangle.$$

Theorem 5 (Global maximum estimate)

Suppose that for some $\alpha_i \in [0, 1]$ and $p_i, q_i \in (1, \infty)$ with $\frac{d}{p_i} + \frac{2}{q_i} < 2 - \alpha_i$, $i = 1, 2, 3$,

$$b \in \tilde{\mathbb{H}}_{q_1}^{-\alpha_1, p_1}, \quad -\operatorname{div} b \leq \Theta_b \in \tilde{\mathbb{H}}_{q_2}^{-\alpha_2, p_2}, \quad f \in \tilde{\mathbb{H}}_{q_3}^{-\alpha_3, p_3}. \quad (4.2)$$

Let $u \in \mathcal{V}_{loc} \cap \mathbb{L}_{loc}^\infty$ be a weak solution of PDE (4.1) with initial value $u(0) = 0$. For any $T > 0$, there exists a constant $C > 0$ depending only on T, d, α_j, p_j, q_j and the quantity

$$\kappa := |||b|||_{-\alpha_1, p_1; q_1} + |||\Theta_b|||_{-\alpha_1, p_1; q_1}$$

such that

$$\|u\|_{L^\infty([0, T] \times \mathbb{R}^d)} + |||u\mathbf{1}_{[0, T]}|||_{\mathcal{V}} \leq C |||f\mathbf{1}_{[0, T]}|||_{-\alpha_3, p_3; q_3}.$$

- When $f \equiv 0$, under (4.2) with $\alpha_j = 0$, the local maximum principle is proved by [Nazarov and Ural'tseva \(2012\)](#) by using Moser's iteration.

- When $f \equiv 0$, under (4.2) with $\alpha_j = 0$, the local maximum principle is proved by [Nazarov and Ural'tseva \(2012\)](#) by using Moser's iteration.
- In elliptic case with $b = 0$ and $f \in L^p(\mathbb{R}^d)$ for $p > \frac{d}{2}$, [Han-Lin \(2011\)](#) show the same maximum principle by De-Giorgi and Moser's iterations.

Thank you for your kind attention!