

# 北京师范大学 随机数学研究中心 学术报告

**题目:** On the topological boundary of the range of super-Brownian motion

**报告人:** 洪杰梁 博士 (*University of British Columbia*)

**时间:** 2021年5月17日, 上午 10:00-11:00

**地点:** 腾讯会议 964 723 428

**摘要:** Let  $\partial\mathcal{R}$  be the topological boundary of the range of super-Brownian motion and  $\dim$  denotes Hausdorff dimension, then with probability one, for any open set  $U$ ,  $U \cap \partial\mathcal{R} \neq \emptyset$  implies

$$\dim(U \cap \partial\mathcal{R}) = d_f := \begin{cases} 4 - 2\sqrt{2} \approx 1.17 & \text{if } d = 2, \\ \frac{9 - \sqrt{17}}{2} \approx 2.44 & \text{if } d = 3. \end{cases}$$

These dimension estimates are also refined in a number of ways. In  $d = 2$  and  $d = 3$ , we construct a random measure  $\mathcal{L}$ , called the boundary local time measure, whose support equals  $\partial\mathcal{R}$ . It is constructed as a rescaled limit of the total local time  $L_\infty^x$  where mass becomes concentrated at points  $x$  where  $L_\infty^x$  is small but positive. It is conjectured that the  $d_f$ -dimensional Minkowski content of  $\partial\mathcal{R}$  is equal to the total mass of the boundary local time  $\mathcal{L}$  up to some constant.