

## MULTI-DIMENSIONAL Q-PROCESSES\*

YAN SHIJIAN (严士健) CHEN MMFA (陈木法)\*\*

### Abstract

In this paper, the authors propose a method which reduces the multi-dimensional problem to one-dimensional ones. By keeping the idea in mind, some sufficient conditions which are much more practical for the uniqueness, recurrency and ergodicity of multi-dimensional Q-processes are obtained.

The conditions are effective not only for the models in non-equilibrium systems, but also for their couplings and others.

### § 1. Introduction

Some stochastic models for linear Master equations of several variables have been introduced in the studies of non-equilibrium systems<sup>[1,2,3]</sup>. In probability language, the models correspond to some Q-processes which satisfy the forward Kolmogorov equation. Thus, one would like to know the uniqueness, the recurrency, and the ergodicity for the Q-processes. It is known that there are some general results about the above problems (cf. [4]). But these results are not effective to those models studied in [3]. As we know, there is only one paper<sup>[5]</sup> which studies directly the properties mentioned above for two-dimensional Q-processes. In this paper, we will propose a method which reduces the multi-dimensional problems to one-dimensional ones. By keeping the idea in mind, we obtain some sufficient conditions which are much more practical for the uniqueness, recurrency and ergodicity of multi-dimensional Q-processes. These conditions are effective not only for the models in [3], but also for their couplings and others.

Now, we are going to state the main results in this paper.

Let  $E$  be a countable set and  $(q(\eta, \zeta): \eta, \zeta \in E)$  be a Q-matrix on  $E \times E$ . Throughout the paper, we will assume that the Q-matrices are totally stable and conservative. Let  $\{E_k \subset E: E_k \neq \phi, k \geq 0\}$  be a disjoint countable partition of  $E$ , and put

---

Manuscript received December 31, 1983.

\* Research supported in part by the Foundation of Zhongshan University Advanced Research Centre.

\*\* Department of Mathematics, Beijing Normal University, Beijing, China.

$$q_{kj} \equiv \begin{cases} \sup\{\sum_{\zeta \in E_j} q(\eta, \zeta) : \eta \in E_k\}, & j > k, \\ \inf\{\sum_{\zeta \in E_j} q(\eta, \zeta) : \eta \in E_k\}, & j < k. \end{cases} \quad (1.1)$$

We say that a  $Q$ -matrix is regular if it determines at most one  $Q$ -process.

The following two results are on the uniqueness for  $Q$ -processes.

**Theorem 1.** Suppose that  $(q(\eta, \zeta) : \eta, \zeta \in E)$  satisfies the following two conditions:

$$q(\eta, \zeta) > 0, \eta \in E_k \Rightarrow \zeta \in \bigcup_{i=0}^{k+1} E_i, \quad (1.2)$$

$$O_k = \sup\{q(\eta) : \eta \in E_k\} < \infty, \quad k \geq 0. \quad (1.3)$$

Then,  $(q(n, \zeta))$  is regular if so is  $(q_{ij})$ .

**Theorem 2.** For each  $k \in Z_+ = \{0, 1, 2, \dots\}$ , let  $B_k$  be a non-empty subset of  $E_k$ . Suppose that  $(q(\eta, \zeta))$  satisfies (1.2), (1.3) and the following conditions:

$$\eta \in E_k, \zeta \in E_{k+1}, q(\eta, \zeta) > 0 \Rightarrow \zeta \in B_{k+1}, \quad (1.4)$$

$$k \in Z_+, \eta \in E_k \setminus B_k, q(\eta, \zeta) > 0 \Rightarrow \zeta \in B_k \cup B_{k+1}. \quad (1.5)$$

Define

$$\tilde{q}_{kj} = \begin{cases} \inf\left\{\sum_{\zeta \in B_j} \left[q(\eta, \zeta) + \sum_{\xi \in B_j \setminus B_k} \frac{q(\eta, \zeta)q(\xi, \zeta)}{q(\xi)}\right] : \eta \in B_k\right\}, & j < k, \\ \sup\left\{\sum_{\zeta \in B_{k+1}} \left[q(\eta, \zeta) + \sum_{\xi \in B_k \setminus B_k} \frac{q(\eta, \xi)q(\xi, \zeta)}{q(\xi)}\right] : \eta \in B_k\right\}, & j = k+1, \\ 0, & j > k+1. \end{cases} \quad (1.6)$$

Then,  $(q(n, \zeta))$  is regular if so is  $(\tilde{q}_{kj})$ .

It is clear that Theorem 2 reduces to Theorem 1 in the case of  $B_k = E_k (k \in Z_+)$ .

Let  $S$  be a finite or countable set, and set  $X = Z_+^S$ . For  $\eta = \{\eta_u : u \in S\} \in X$ , we put

$$|\eta| = \sum_{u \in S} \eta_u, \quad X_0 = \{\eta \in X : |\eta| < \infty\}.$$

Clearly,  $X_0 = X$  if  $S$  is finite. We use  $\theta$  to denote the element in  $X_0$  so that  $|\theta| = 0$ .

If  $E = X_0$  in Theorems 1 and 2, we see that  $(q(\eta, \zeta))$  is a multi-dimensional (even infinite dimensional)  $Q$ -matrix. What the above theorems mean is reducing the uniqueness problem in multi-dimensions to the one in one-dimension by choosing an appropriate partition  $\{E_k\}_0^\infty$  of  $E$ . This idea is very useful since the  $Q$ -matrix  $(q_{ij})$  ( $(\tilde{q}_{ij})$ ) in (1.1) ((1.6)) is a generalized birth-death  $Q$ -matrix, for which we have the following uniqueness criterion.

**Theorem 3.**

(i) Suppose that

$$q_{kj} = 0, j > k+1; q_{k, k+1} > 0, k, j \in Z_+. \quad (1.7)$$

Then  $(q_{kj})$  is regular iff

$$R \equiv \sum_{k=0}^{\infty} m_k = \infty, \quad (1.8)$$

where

$$\left\{ \begin{aligned} m_k &\equiv \sum_{i=0}^k F_k^{(i)} / q_{i,i+1}, \quad k \in Z_+, \\ F_k^{(k)} &\equiv 1, \quad k \in Z_+, \\ F_k^{(i)} &\equiv q_{k,k+1}^{-1} \sum_{j=i}^{k-1} q_k^{(j)} F_j^{(i)}, \quad 0 \leq i < k, \\ q_k^{(i)} &= \sum_{j=0}^i q_{kj}, \quad 0 \leq i < k. \end{aligned} \right. \quad (1.9)$$

(ii) Suppose that (1.7) holds except  $q_{01} = 0$ . Then  $(q_{ij})$  is regular iff


$$R' = \infty, \quad (1.10)$$

where  $R'$  can be obtained from (1.8) and (1.9) when 0 is replaced by 1 and  $Z_+$  is replaced by  $Z_+ \setminus \{0\}$ .

The above three theorems will be proved in § 2. Theorem 3 is an extension of [6, § 3, Corollary 1], for which our proof is much more simple.

It is clear that both  $R$  and  $R'$  are computable, so they are very convenient in the practice. As their applications, in § 3, we will discuss the uniqueness for the following models<sup>[3]</sup>:

*An autocatalytic production of a chemical*




Its  $Q$ -matrix is

$$q(\eta, \zeta) = \begin{cases} \lambda_1 a_u \eta_u, & \zeta = \eta + e_u, \\ \lambda_2 \binom{\eta_u}{2}, & \zeta = \eta - 2e_u, \\ \eta_u p(u, v), & \zeta = \eta - e_u + e_v, \quad u \neq v, \\ 0, & \text{other } \zeta \neq \eta, \end{cases} \quad \eta, \zeta \in X_0, \quad (1.11)$$

where  $S$  is the set of seats,  $u, v \in S$ ,  $e_u = \{\delta_{uv} : v \in S\}$ ,  $a_u$  and  $\eta_u (u \in S)$  are the numbers of  $A$ -particles and  $X$ -particles, respectively,  $p(u, v)$  is the transition rate of an  $X$ -particle from  $u$  to  $v$ . We will assume that

$$\sum_v p(u, v) \leq C < \infty, \quad u \in S$$


*Schlögl model:*



Its  $Q$ -matrix is

$$q(\eta, \zeta) = \begin{cases} \lambda_1 a_u \binom{\eta_u}{2} + \lambda_4 b_u, & \zeta = \eta + e_u, \\ \lambda_2 \binom{\eta_u}{3} + \lambda_3 \eta_u, & \zeta = \eta - e_u, \\ \eta_u p(u, v), & \zeta = \eta - e_u + e_v, \quad u \neq v, \\ 0, & \text{other } \zeta \neq \eta. \end{cases} \quad (1.12)$$

Lotka-Volterra model:




Its state space should be  $X = (Z_+^2)^S$ . For each  $\eta \in X$ ,  $n = \{\eta_{ui} : u \in S, i = 1, 2\}$ , set  $|\eta| = \sum_{u \in S} \sum_{i=1}^2 n_{ui}$ . Then  $X_0 = \{n \in X : |n| < \infty\}$ .  $n_{ui}$  denotes the numbers of  $X_i$ -particles in  $u$ , and we use  $p_i(u, v)$  to denote the transition rate of an  $X_i$ -particle from  $u$  to  $v$ . Now, the Q-matrix for the model can be written as follows:

$$q(\eta, \zeta) = \begin{cases} \lambda_1 a_u \eta_{u1}, & \zeta = \eta + e_{u1}, \\ \lambda_3 b_u \eta_{u2}, & \zeta = \eta - e_{u2}, \\ \lambda_2 \eta_{u1} \eta_{u2}, & \zeta = \eta - e_{u1} + e_{u2}, \\ \eta_{ui} p_i(u, v), & \zeta = \eta - e_{ui} + e_{vi}, \quad i = 1, 2, u \neq v, \\ 0, & \text{other } \zeta \neq \eta, \end{cases} \quad (1.13)$$

where  $e_{ui} = \{\delta_{uv} \delta_{ij} : v \in S, j = 1, 2\}$ .

Brusselator:



Again, the state space is  $X = (Z_+^2)^S$ . Its Q-matrix is

$$q(\eta, \zeta) = \begin{cases} \lambda_1 a_u, & \zeta = \eta + e_{u1}, \\ \lambda_4 \eta_{u1}, & \zeta = \eta - e_{u1}, \\ \lambda_2 b_u \eta_{u1}, & \zeta = \eta - e_{u1} + e_{u2}, \\ \lambda_3 \binom{\eta_{u1}}{2} \eta_{u2}, & \zeta = \eta - e_{u2} + e_{u1}, \\ \eta_{ui} p_i(u, v), & \zeta = \eta - e_{ui} + e_{vi}, \quad i = 1, 2, u \neq v, \\ 0, & \text{other } \zeta \neq \eta. \end{cases} \quad (1.14)$$

The point of applying our results to the above models is simply choosing

$$E_k = \{\eta \in X_0 : |\eta| = k\}.$$

As another application of Theorem 1 and Theorem 3, we will discuss the uniqueness for coupling Q-processes. To state this, notice that there is a one-to-one mapping between Q-matrix  $(q(\eta, \zeta))$  and the following operator:

$$\Omega f(\eta) = \sum_{\zeta \in X_0} q(\eta, \zeta) (f(\zeta) - f(\eta)), \quad \eta \in X_0, \quad (1.15)$$

where  $f$  is a bounded real function on  $X_0$ . So we can also use the term "Q-process" instead of Q-process.

**Theorem 5.** Let  $S$  be finite,  $X = Z_+^S$  and

$$(\Omega f)(\eta) = \sum_u \beta(u, \eta_u) (f(\eta + e_u) - f(\eta)) + \sum_u \delta(u, \eta_u) (f(\eta - e_u) - f(\eta)) \\ + \sum_{u,v} \gamma(u, v, \eta_u, \eta_v) (f(\eta - e_u + e_v) - f(\eta)), \quad (1.16)$$

where  $\beta, \delta$  and  $\gamma$  are non-negative and  $\delta(u, 0) = 0$ . Suppose that  $\beta(u, k)$  and  $\gamma(u, v, k, l)$  are increasing in  $k$  and  $l$ , and define the coupling operator  $\tilde{Q}$  as follows:

$$\begin{aligned} \tilde{Q}f(\eta, \zeta) = & \sum_{u: \eta_u = \zeta_u} \{ \beta(u, \eta_u) [f(\eta + e_u, \zeta) - f(\eta, \zeta)] + \delta(u, \eta_u) [f(\eta - e_u, \zeta) - f(\eta, \zeta)] \\ & + \beta(u, \zeta_u) [f(\eta, \zeta + e_u) - f(\eta, \zeta)] + \delta(u, \zeta_u) [f(\eta, \zeta - e_u) - f(\eta, \zeta)] \\ & + \sum_{u: \zeta_u = \eta_u} \{ \beta(u, \eta_u) \wedge \beta(u, \zeta_u) [f(\eta + e_u, \zeta + e_u) - f(\eta, \zeta)] \\ & + \delta(u, \eta_u) \wedge \delta(u, \zeta_u) [f(\eta - e_u, \zeta - e_u) - f(\eta, \zeta)] \} \\ & + \sum_{u, v} (\gamma(u, v, \eta_u, \eta_v) - \gamma(u, v, \zeta_u, \zeta_v))^+ \\ & \cdot [f(\eta - e_u + e_v, \zeta) - f(\eta, \zeta)] \\ & + \sum_{u, v} (\gamma(u, v, \zeta_u, \zeta_v) - \gamma(u, v, \eta_u, \eta_v))^+ \\ & \cdot [f(\eta, \zeta - e_u + e_v) - f(\eta, \zeta)] \\ & + \sum_{u, v} \gamma(u, v, \eta_u, \eta_v) \wedge \gamma(u, v, \zeta_u, \zeta_v) \\ & \cdot [f(\eta - e_u + e_v, \zeta - e_u + e_v) - f(\eta, \zeta)], \end{aligned}$$

where  $f$  is a bounded real function on  $X \times X$ . Set

$$\left\{ \begin{aligned} r_k &= \max \left\{ \sum_u \beta(u, \eta_u) : |\eta| = k \right\}, \\ s_k &= \min \left\{ \sum_u \delta(u, \eta_u) : |\eta| = k \right\}, k \in Z_+, \\ \bar{R} &= \begin{cases} \sum_{k=1}^{\infty} \left[ \frac{1}{2r_k} + \dots + \frac{s_k \cdots s_2}{2^k r_k \cdots r_1} \right], r_0 = 0, r_k > 0, k \geq 1, \\ \sum_{k=0}^{\infty} \left[ \frac{1}{2r_k} + \dots + \frac{s_k \cdots s_1}{2^{k+1} r_k \cdots r_0} \right], r_k > 0, k \in Z_+. \end{cases} \end{aligned} \right. \quad (1.18)$$

If

$$\bar{R} = +\infty, \quad (1.19)$$

then, both  $\Omega$ -process and  $\tilde{Q}$ -process are unique. Moreover

$$\sum_{\xi \in X} \tilde{p}(t, (\eta, \zeta), (\tilde{\eta}, \tilde{\xi})) = p(t, \eta, \tilde{\eta}) \quad (1.20)$$

$$\sum_{\tilde{\eta} \in X} \tilde{p}(t, (\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) = p(t, \zeta, \tilde{\zeta}), \quad (1.21)$$

and for each  $\eta \leq \zeta$  (i.e.  $\eta_u \leq \zeta_u, u \in S$ ), we have

where  $p(t, \eta, \zeta)$  ( $p(t, (\eta, \zeta), (\tilde{\eta}, \tilde{\zeta}))$ ) is the  $\Omega$ -process ( $\tilde{Q}$ -process).

Of course, the approach used to prove Theorem 5 can be generalized, but we will not do it in this paper. It is easy to check (see § 4) that the  $Q$ -matrices defined in (1.12), (1.13) and (1.14) satisfy the assumptions of Theorem 5. Hence, this theorem works for the above models.

The recurrence for  $Q$ -processes will be studied in § 5. The main result is the following

**Theorem 6.** Let  $E = X_0$  and let  $(q(\eta, \zeta))$  satisfy (1.2) and (1.3). Let  $E_0 = \{\theta\}$ ,  $E_k$  ( $k \geq 1$ ) be finite. Suppose that the  $Q$ -matrices  $(q(\eta, \zeta))$  and  $(q_i)$  defined by (1.1)

is irreducible and regular. Then, the  $(q(\eta, \zeta))$ -process is recurrent if so is the  $(q_{ij})$ -process. Moreover, the  $(q_{ij})$ -process is recurrent iff

$$\sum_{k=0}^{\infty} F_k^{(0)} = \infty,$$

where  $F_k^{(0)}$  is defined in (1.9).

The positive recurrency and ergodicity for  $(q(\eta, \zeta))$ -processes will be studied in § 6. The main results are the following

**Theorem 7.** Let  $(q_{ij})$  be an arbitrary irreducible and regular Q-matrix on  $Z_+^2$ . Then the  $(q_{ij})$ -process is positive recurrent iff there exists a non-negative solution  $(x_i; i \in Z_+)$  to the following inequalities:

$$\sum_j q_{ij} x_j + 1 \leq 0, \quad i \neq i_0, \quad \left| \sum_j q_{i_0, j} x_j \right| < \infty, \quad (1.22)$$

for some  $i_0 \in Z_+$  (equivalently, for any  $i_0 \in Z_+$ ).

**Theorem 8.** Let  $S$  be finite, and let  $(q(\eta, \zeta))$  satisfy (1.2) and (1.3) with  $E_0 = \{\theta\}$ . Suppose that both  $(q(\eta, \zeta))$  and  $(q_{ij})$  defined by (1.1) are irreducible. If  $(q_{ij})$  is regular and for (1.22), there exists a nonnegative solution  $(u_i)$  for  $i_0 = 0$ , which is increasing in  $i$ , then  $(q(\eta, \zeta))$ -process is positive recurrent. In fact, it is ergodic, i.e. there exists a probability measure  $\{\mu(\eta)\}$  on  $X$  such that

$$\lim_{t \rightarrow \infty} p(t, \xi, \eta) = \mu(\eta), \quad \xi, \eta \in X, \quad (1.23)$$

where  $p(t, \xi, \eta)$  is the Q-process corresponding to  $(q(\eta, \zeta))$ .

**Theorem 9.** Let  $(q_{ij})$  be a Q-matrix satisfying (1.7). Then, there exists a non-negative and increasing solution  $(x_i)$  to (1.22) with  $i_0 = 0$ , iff

$$d \equiv \sup_{k \in Z_+} d_k / F_k^{(0)} < \infty, \quad (1.24)$$

where  $F_k^{(0)}$  is defined by (1.9) and

$$d_0 = 0, \quad d_k = q_{k, k+1}^{-1} \left( 1 + \sum_{s=0}^{k-1} q_k^{(s)} d_s \right), \quad k > 0. \quad (1.25)$$

Moreover, if (1.24) holds, then the function  $(u_i; i \in Z_+)$  defined by

$$u_0 = 0, \quad u_1 \geq d, \quad u_{k+1} = u_k + F_k^{(0)} u_1 - d_k, \quad k \geq 1 \quad (1.26)$$

is a non-negative and increasing solution to (1.22) with  $i_0 = 0$ .

As an application, we will show in § 6 that Schlögl model is ergodic. For the models defined in (1.11) and (1.13), we have nothing to do since  $\theta$  is an absorbing state for these models. However, for the Brusselator model defined in (1.14), our conditions do not work, this is a remainder problem.

One may ask whether the condition (1.24) is equivalent to the positive recurrence for the  $(q_{ij})$ -process or not. The answer is negative. To compare the two properties, we have

**Theorem 10.** Let  $(q_{ij})$  be a Q-matrix satisfying (1.7). Then there exists a non-negative solution to (1.22) with  $i_0 = 0$  iff

$$\hat{d} \equiv \sup \left\{ \left( \sum_{i=0}^k d_i \right) / \left( \sum_{i=0}^k F_i^{(0)} \right) : k \in Z_+ \right\} < \infty. \quad (1.27)$$

If the  $Q$ -matrix  $(q_{ij})$  is also irreducible and regular, then the condition (1.27) is equivalent to the positive recurrence of the  $(q_{ij})$ -process.

It is obvious that  $\hat{d} \leq d$ . For birth-death processes, it is not difficult to show that  $d < \infty$  iff  $\hat{d} < \infty$ .

## § 2. Uniqueness

**Lemma 1.** Let  $(q_{ij}) : i, j \in E$  be a  $Q$ -matrix. Then

$$(\lambda + q_i)u_i \leq \sum_{j \neq i} q_{ij}u_j, \quad 0 \leq u_i \leq 1, \quad i \in E, \lambda > 0 \quad (2.1)$$

has only zero solution iff

$$(\lambda + q_i)u_i = \sum_{j \neq i} q_{ij}u_j, \quad 0 \leq u_i \leq 1, \quad i \in E, \lambda > 0 \quad (2.2)$$

has only zero solution.

*Proof* It is well known that the maximal solution  $(u_i^*)$  to (2.2) can be obtained by the following procedure: define

$$u_i^{(0)} \equiv 1, \quad i \in E, \\ u_i^{(n+1)} = \sum_{j \neq i} q_{ij}u_j^{(n)} / (\lambda + q_i), \quad n \geq 0, i \in E,$$

then

$$u_i^{(n)} \searrow u_i^* \text{ as } n \rightarrow \infty \text{ for each } i \in E.$$

Now, suppose that  $(v_i)$  is a non-zero solution to (2.1). By induction, it is easy to show that  $v_i \leq u_i^{(n)}$  for each  $n \geq 1$  and  $i \in E$ . Hence  $u_i^* \geq v_i$  ( $i \in E$ ). This is a contradiction.

*Proof of Theorem 1.*

By [8, § 4.3, Corollary 1] or [9, § 5.4, Theorem 1], it suffices to prove that for some  $\lambda > 0$  (or equivalently, for each  $\lambda > 0$ ),

$$(\lambda + q(\eta))u(\eta) = \sum_{\zeta \neq \eta} q(\eta, \zeta)u(\zeta), \quad 0 \leq u(\zeta) \leq 1, \eta \in E \quad (2.3)$$

has only zero solution. Suppose that there exists a non-zero solution  $\{u(\eta) : \eta \in E\}$  for some  $\lambda > 0$ . Set

$$u_k = \sup \{u(\eta) : \eta \in E_k\}, \quad k \in Z_+. \quad (2.4)$$

Then  $(u_k : k \in Z_+)$  is non-zero. For each  $k \in Z_+$ , choose  $\varepsilon_k > 0$  and  $\eta^{(k)} \in E_k$  so that

$$\varepsilon_k(\lambda + c_k) < \frac{\lambda}{2} \text{ and } u(\eta^{(k)}) \geq (1 - \varepsilon_k)u_k. \quad (2.5)$$

Replacing  $\eta$  in (2.3) with  $\eta^{(k)}$ , it follows from (1.2), (1.3), (2.4) and (2.5) that

$$\frac{\lambda}{2} u_k + \left[ \sum_{j=0}^{k-1} \sum_{\zeta \in E_j} q(\eta^{(k)}, \zeta) + \sum_{\eta^{(k)} \neq \zeta \in E_k} q(\eta^{(k)}, \zeta) + \sum_{\zeta \in E_{k+1}} q(\eta^{(k)}, \zeta) \right] u_k \\ \leq u_k(\lambda - \varepsilon_k(\lambda + c_k)) + q(\eta^{(k)})u_k \leq (\lambda + q(\eta^{(k)}))(1 - \varepsilon_k)u_k$$

$$\begin{aligned} &\leq (\lambda + q(\eta^{(k)}))u(\eta^{(k)}) = \sum_{\zeta \neq \eta} q(\eta^{(k)}, \zeta)u(\zeta) \\ &\leq \sum_{j=0}^{k-1} \sum_{\zeta \in E_j} q(\eta^{(k)}, \zeta)u_j + \sum_{\eta^{(k)} \neq \zeta \in E_k} q(\eta^{(k)}, \zeta)u_k + \sum_{\zeta \in E_{k+1}} q(\eta^{(k)}, \zeta)u_{k+1}, \end{aligned}$$

i.e.,

$$\frac{\lambda}{2}u_k + \sum_{j=0}^{k-1} \left( \sum_{\zeta \in E_j} q(\eta^{(k)}, \zeta) \right) (u_k - u_j) \leq \sum_{\zeta \in E_{k+1}} q(\eta^{(k)}, \zeta) (u_{k+1} - u_k). \quad (2.6)$$

Clearly,  $u_k$  is increasing. By (1.1), we now get

$$\frac{\lambda}{2}u_k + \sum_{j=0}^{k-1} q_{kj}(u_k - u_j) \leq q_{k, k+1}(u_{k+1} - u_k), \quad k \in Z_+. \quad (2.7)$$

By (1.1) and (1.2), this means that

$$\left( \frac{\lambda}{2} + q_k \right) u_k \leq \sum_{j \neq k} q_{kj} u_j, \quad 0 \leq u_k \leq 1, \quad k \in Z_+. \quad (2.8)$$

But this is impossible by Lemma 1 and the regularity for  $(q_{ij})$ .

*Proof of Theorem 2.* The proof is similar to the proof of Theorem 1. We leave it to the reader as an exercise.

*Proof of Theorem 3.* (1). Suppose that  $(q_{ij})$  satisfies (1.7). Then there exists only one increasing solution to

$$(\lambda + q_i)u_i = \sum_{j \neq i} q_{ij}u_j, \quad u_0 = 1, \quad i \in Z_+ \quad (2.9)$$

for each  $\lambda > 0$ . In fact, we have

$$u_0 = 1, \quad (2.10)$$

$$u_{i+1} = \left[ (\lambda + q_i)u_i - \sum_{j=0}^{i-1} q_{ij}u_j \right] / q_{i, i+1}, \quad i \geq 0,$$

$$u_{i+1} - u_i = \left[ \sum_{j=0}^{i-1} q_{ij}(u_i - u_j) + \lambda u_i \right] / q_{i, i+1}, \quad i \geq 0. \quad (2.11)$$

We now prove that

$$\lambda u_0 m_k \leq u_{k+1} - u_k \leq (u_1 - u_0) F_k^{(0)} + \lambda u_k m_k, \quad k \in Z_+. \quad (2.12)$$

When  $k=1$ , (2.12) holds since

$$u_1 - u_0 = \lambda u_0 / q_{01} = \lambda u_0 m_0.$$

Suppose that it holds for  $k < n$ . Then, by using

$$\sum_{i=0}^{k-1} q_{ki}(u_k - u_i) = \sum_{i=0}^{k-1} q_k^{(i)}(u_{i+1} - u_i), \quad k \in Z_+, \quad (2.13)$$

(2.11), (1.9) and the increasing property of  $u_k$ , it follows that

$$\begin{aligned} u_{n+1} - u_n &= \frac{1}{q_{n, n+1}} \left[ \sum_{k=0}^{n-1} q_n^{(k)}(u_{k+1} - u_k) + \lambda u_n \right] \\ &\leq q_{n, n+1}^{-1} \left[ (u_1 - u_0) \sum_{k=0}^{n-1} q_n^{(k)} F_k^{(0)} + \lambda u_n \left( \sum_{k=0}^{n-1} q_n^{(k)} m_k + 1 \right) \right] \\ &= (u_1 - u_0) F_n^{(0)} + \lambda u_n \left[ \sum_{j=0}^{n-1} q_{j, j+1}^{-1} q_{n, n+1}^{-1} \sum_{k=j}^{n-1} q_n^{(k)} F_k^{(j)} + q_{n, n+1}^{-1} F_n^{(n)} \right] \\ &= (u_1 - u_0) F_n^{(0)} + \lambda u_n m_n, \\ u_{n+1} - u_n &\geq q_{n, n+1}^{-1} \left[ \sum_{k=0}^{n-1} \lambda u_0 q_n^{(k)} m_k + \lambda u_n \right] \geq \lambda u_0 q_{n, n+1}^{-1} \left( \sum_{k=0}^{n-1} q_n^{(k)} m_k + 1 \right) = \lambda u_0 m_n. \end{aligned}$$



By induction, this proves (2.12).

(2). To prove (i) of Theorem 3, it is enough to show that (2.2) has only zero solution iff (1.8) holds. Suppose that  $R < \infty$  and  $(u_k)$  is the solution to (2.9) constructed by (2.10). By (2.12) and (1.9), we get

$$u_{k+1}u_k^{-1} - 1 \leq (u_1 - u_0)u_k^{-1}F_k^{(0)} + \lambda m_k \leq [\lambda + (u_1 - u_0)q_{01}]m_k.$$

Hence  $u_{k+1}u_k^{-1} - 1 < 1/2$ , and so  $\log(u_{k+1}u_k^{-1}) \leq 2(u_{k+1}u_k^{-1} - 1)$  for  $k$  large enough. Therefore, there exists a constant  $C > 0$  such that

$$\log u_k = \sum_{i=1}^{k-1} \log(u_{i+1}u_i^{-1}) \leq CR < \infty.$$

This implies that

$$u_\infty = \lim_{k \rightarrow \infty} u_k < \infty.$$

Now,  $\hat{u}_k \equiv u_k u_\infty^{-1} \in [0, 1]$  is a non-zero solution to (2.2). Conversely, if (2.2) has a non-zero solution, it is easy to show that  $R < \infty$ .

As for (ii) of Theorem 3, it is enough to notice that (2.2) has only zero solution iff

$$(\lambda + q_k)u_k = \sum_{j \neq 0, k} q_{kj}u_j, \quad 0 \leq u_k \leq 1, \quad k \geq 1$$

has only zero solution in the case of  $q_{01} = 0$ .

### § 3. Uniqueness for Some Processes of Non-equilibrium Systems

As applications of the results in § 1, we will prove in this section that the  $Q$ -matrices defined by (1.11)–(1.14) are regular, i. e. they determine uniquely the  $Q$ -processes.

#### Theorem 4.

(i) The  $Q$ -matrix  $(q(\eta, \zeta): \eta, \zeta \in X_0)$  defined in (1.11) is regular if

$$a \equiv \sup\{a_u: u \in S\} < \infty; \quad (3.1)$$

(ii) Let  $S$  be finite, then the  $Q$ -matrix  $(q(\eta, \zeta): \eta, \zeta \in X)$  defined by (1.12) is regular;

(iii) The  $Q$ -matrix  $(q(\eta, \zeta): \eta, \zeta \in X_0)$  defined by (1.13) is regular if (3.1) holds;

(iv) The  $Q$ -matrix  $(q(\eta, \zeta): \eta, \zeta \in X_0)$  defined by (1.14) is regular if

$$\tilde{a} \equiv \sum_{u \in S} a_u < \infty. \quad (3.2)$$

*Proof* Take  $E_k = \{\eta \in X_0: |\eta| = k\}$ ,  $k \in \mathbb{Z}_+$  in the four cases. Then (1.2) and (1.3) hold.

For (i) and (iii), the conditions of Theorem 3. (ii) hold, and

$$q_{k, k+1} \leq \lambda_1 a k.$$

Since

$$R' = \sum_{k=1}^{\infty} \left( \sum_{i=1}^{\infty} F_k^{(i)} / q_{i,i+1} \right) \geq \sum_{k=1}^{\infty} F_k^{(k)} q_{k,k+1}^{-1} \geq \sum_{k=1}^{\infty} (\lambda_1 a k)^{-1} = \infty,$$

now, (i) and (iii) follow from Theorem 3. (ii).

The  $Q$ -matrices  $(q_{ij})$  corresponding to (1.12) and (1.14) are birth-death  $Q$ -matrices. By (1.8) and (1.9), it is easy to carry out that

$$R = \sum_{k=0}^{\infty} \sum_{i=0}^k \frac{s_k \cdots s_{i+1}}{r_k \cdots r_{i+1} r_i}, \tag{3.3}$$

$s_k \equiv q_{k,k-1}, r_k \equiv q_{k,k+1}.$

For (ii), we have

$$r_k = \sup \left\{ \sum_{u \in S} \left[ \lambda_1 a_u \binom{\eta_u}{2} + \lambda_4 b_u \right] : |\eta| = k \right\} \leq 2^{-1} \lambda_1 a (k^2 - k) + \lambda_4 \sum_{u \in S} b_u. \tag{3.4}$$

Here we have used

$$\sup \left\{ \sum_{u \in S} \eta_u^2 : |\eta| = k \right\} = k^2. \tag{3.5}$$

By (3.5) and

$$\left( \sum_u \eta_u / |S| \right)^2 \leq \sum_u \eta_u^2 / |S|$$

we see that

$$\begin{aligned} s_k &= \inf \left\{ \sum_{u \in S} \left[ \lambda_2 \binom{\eta_u}{3} + \lambda_3 \eta_u \right] : |\eta| = k \right\} \\ &\geq 6^{-1} \lambda_2 \left[ \inf_{|\eta|=k} \sum_{u \in S} \eta_u^3 - 3 \sup_{|\eta|=k} \sum_{u \in S} \eta_u^2 \right] + (\lambda_3 + \lambda_2/3) k \\ &> \frac{\lambda_2}{6} \left[ |S| \left( \frac{k}{|S|} \right)^3 - 3k^2 \right] + (\lambda_3 + \lambda_2/3) k. \end{aligned}$$

Now, by (3.4) and (3.6), we get

$$\lim_{k \rightarrow \infty} \sum_{i=0}^k \frac{s_k \cdots s_{i+1}}{r_k \cdots r_{i+1} r_i} \geq \lim_{k \rightarrow \infty} \frac{s_k s_{k-1}}{r_k r_{k-1} r_{k-2}} \geq 2\lambda_1^2 / (9|S|^4 (\lambda_1 a)^3).$$

It follows from (3.3) that  $R = \infty$ . Therefore (ii) follows from Theorem 3. (i).

For (iv), we have

$$r_k = \sup \left\{ \sum_u \lambda_1 a_u : |\eta| = k \right\} = \lambda_1 \tilde{a}, k \in Z_+.$$

By (3.3), we get

$$R \geq \sum_{k=0}^{\infty} \frac{1}{r_k} = \infty.$$

Therefore (iv) follows also from Theorem 3. (i).

To conclude this section, we discuss two special cases.

**Proposition 1.** Let  $(\bar{q}_{ij} : i, j \in Z_+)$  be a  $Q$ -matrix satisfying

$$\bar{q}_{ij} = 0, |i - j| > 2 \tag{3.6}$$

and

$$\begin{cases} r_k \equiv \max(\bar{q}_{2k, 2k+2}, \bar{q}_{2k+1, 2k+2} + \bar{q}_{2k+1, 2k+3}) > 0, \\ s_k \equiv \min(\bar{q}_{2k, 2k-2} + \bar{q}_{2k, 2k-1}, \bar{q}_{2k+1, 2k-1}) > 0, \end{cases} \tag{3.7}$$

$$\sum_{k=0}^{\infty} \sum_{i=0}^k \frac{s_k \cdots s_{i+1}}{r_k \cdots r_{i+1} r_i} = \infty. \tag{3.8}$$

Then  $(\bar{q}_{ij})$  is regular.

*Proof* Simply take  $E_k = \{2k, 2k+1\}$ ,  $k \in Z_+$  and apply Theorems 1 and 3.

**Proposition 2.** Let  $(\bar{q}_{ij}: i, j \in Z_+)$  be a  $Q$ -matrix satisfying

$$\begin{aligned} \bar{q}_{ij} &= 0, \quad i, j \geq n \text{ and } |i-j| > 1, \\ \bar{q}_{n+k, n+k+1} &> 0, \quad k \in Z_+, \end{aligned} \tag{3.9}$$

where  $n$  is a given positive integer. Set  $r_k = \bar{q}_{n+k, n+k+1}$ ,  $k \geq 0$  and  $s_k = \bar{q}_{n+k, n+k-1}$ ,  $k \geq 1$ . Then  $(\bar{q}_{ij})$  is regular if (3.8) holds.

*Proof* Simply take  $E_0 = \{0, 1, 2, \dots, n\}$ ,  $E_k = \{n+k\}$ ,  $k \geq 1$  and apply Theorems 1 and 3.

### § 4. Further Applications

In order to prove Theorem 5, we introduce a relation “ $\rightarrow$ ” on  $X \times X$  as follows:

$$(\eta, \zeta) \rightarrow \begin{cases} (\eta + e_u, \zeta) \text{ or } (\eta, \zeta + e_u), \eta_u \neq \zeta_u \\ (\eta - e_u, \zeta), \eta_u \neq \zeta_u, \eta_u \geq 1, \\ (\eta, \zeta - e_u), \eta_u \neq \zeta_u, \zeta_u \geq 1, \\ (\eta + e_u, \zeta + e_u), \eta_u = \zeta_u, \\ (\eta - e_u, \zeta - e_u), \eta_u = \zeta_u \geq 1 \end{cases} \tag{4.1}$$

for each  $u \in S$ . Then, define

$$\begin{cases} \tilde{B}_0 = \{(\theta, \theta)\}, \\ \tilde{B}_{k+1} = \{(\eta, \zeta) : \exists (\tilde{\eta}, \tilde{\zeta}) \in \bigcup_{i=0}^k \tilde{B}_i \text{ such that } (\eta, \zeta) \rightarrow (\tilde{\eta}, \tilde{\zeta})\} \cup \tilde{B}_0, \\ \tilde{E}_0 = \tilde{B}_0, \tilde{E}_{k+1} = \tilde{B}_{k+1} \setminus \tilde{B}_k, k \in Z_+. \end{cases} \tag{4.2}$$

**Lemma 1.**

- (i) For each  $n \in Z_+$ ,  $\tilde{B}_n \subset \tilde{B}_{n+1}$ ;
- (ii) For each  $n \in Z_+$ ,

$$\tilde{B}_n = \{(\theta, \theta)\} \cup \left\{ \left( \sum_{i=1}^{h_m} e_{u_i}, \sum_{i=h_m+1}^m e_{u_i} \right) : \right. \tag{4.3}$$

$$\left. m = 1, 2, \dots, n; h_m = 0, 1, \dots, m; S \ni u_1, \dots, u_n \text{ may be repeated} \right\}$$

$$\cup \left\{ \left( \sum_{i=1}^{k_i} e_{u_i}, \sum_{i=1}^i e_{u_i} + \sum_{i=k_i+1}^n e_{u_i} \right) : i = 1, 2, \dots, n; k_i = i, i+1, \dots, n; \right.$$

$$\left. S \ni u_1, \dots, u_n \text{ may be repeated} \right\},$$

where  $\sum_{i=a}^b e_{u_i} = \theta$  whenever  $a > b$ ;

- (iii)  $\tilde{E}_n (n \in Z_+)$  are disjointed,  $\bigcup_{n=0}^{\infty} \tilde{E}_n = X \times X$  and  $\{\tilde{E}_n\}$  satisfies (1.2) and (1.3).

*Proof* (i) Clearly,  $\tilde{B}_0 \subset \tilde{B}_1$ . For each  $(\eta, \zeta) \in \tilde{B}_n$  ( $n \geq 1$ ), by (4.2), there exists

an  $(\tilde{\eta}, \tilde{\zeta}) \in \bigcup_{i=0}^{n-1} \tilde{B}_i \subset \bigcup_{i=0}^n \tilde{B}_i$  such that  $(\eta, \zeta) \rightarrow (\tilde{\eta}, \tilde{\zeta})$ . Hence  $(\eta, \zeta) \in \tilde{B}_{n+1}$ . By induction this proves (i).

(ii) To prove (4.3), denote the right side of (4.3) by  $\bar{B}_n$ . Clearly  $\tilde{B}_0 = \bar{B}_0$ . Suppose that  $\tilde{B}_n = \bar{B}_n$  for all  $n < k$ . We then have to show that  $\tilde{B}_n = \bar{B}_n$  for  $n = k$ .

First, we consider such an element  $(\eta, \zeta)$  that has the form  $(\sum_{i=1}^{h_m} e_{u_i}, \sum_{i=h_m+1}^m e_{u_i})$ . If  $m < k$ , then

$$(\eta, \zeta) = (\sum_{i=1}^{h_m} e_{u_i}, \sum_{i=h_m+1}^m e_{u_i}) \in \bar{B}_m = \tilde{B}_m \subset \tilde{B}_k.$$

We may now assume that  $m = k$ . Hence

$$(\eta, \zeta) = (\sum_{i=1}^{h_k} e_{u_i}, \sum_{i=h_k+1}^k e_{u_i}).$$

If  $h_k = k$ , then

$$(\eta, \zeta) = (\sum_{i=1}^k e_{u_i}, \theta) \rightarrow (\sum_{i=1}^{k-1} e_{u_i}, \theta) \in \tilde{B}_{k-1}.$$

If  $h_k < k$ , then there are two cases:

(a) The times of  $u_k$  appeared in  $\{u_1, \dots, u_{h_k}\}$  and in  $\{u_{h_k+1}, \dots, u_k\}$  are different.

Then

$$(\eta, \zeta) \rightarrow (\eta, \zeta - e_{u_k}) = (\sum_{i=1}^{h_k} e_{u_i}, \sum_{i=h_k+1}^{k-1} e_{u_i}) \in \tilde{B}_{k-1}.$$

(b) The times of  $u_k$  appeared in  $\{u_1, \dots, u_{h_k}\}$  and in  $\{u_{h_k+1}, u_k\}$  are the same.

Then we may assume  $u_{h_k} = u_k$ . Hence

$$(\eta, \zeta) \rightarrow (\eta - e_{u_k}, \zeta - e_{u_k}) \in \tilde{B}_{k-1}.$$

Therefore, in both cases (a) and (b), we have  $(\eta, \zeta) \in \tilde{B}_k$ .

Next, we consider such an element  $(\eta, \zeta)$  that has the form  $(\sum_{i=1}^{k_1} e_{u_i}, \sum_{i=1}^k e_{u_i} + \sum_{i=k_1+1}^k e_{u_i})$ . There are three cases:

(a) The times of  $u_1$  appeared in  $\{u_1, \dots, u_{k_1}\}$  and in  $\{u_1, \dots, u_i, u_{k_1+1}, \dots, u_k\}$  are the same. Then  $(\eta, \zeta) \rightarrow (\eta - e_{u_1}, \zeta - e_{u_1}) \in \tilde{B}_{k-1}$ .

Now, assume that the times of  $u_1$  appeared in the above two sets are different. Then,  $u_1$  should appear in  $\{u_{k_1+1}, \dots, u_{k_1+1}, \dots, u_k\}$ .

(b) If  $u_1 \in \{u_{k_1+1}, \dots, u_{k_1}\}$ , we may assume that  $u_{k_1} = u_1$ . Then

$$(\eta, \zeta) \rightarrow (\eta - e_{u_1}, \zeta) \in \tilde{B}_{k-1}.$$

(c) If  $u_1 \in \{u_{k_1+1}, \dots, u_k\}$ , then

$$(\eta, \zeta) \rightarrow (\eta, \zeta - e_{u_1}) \in \tilde{B}_{k-1}.$$

In the above three cases, we always have  $(\eta, \zeta) \in \tilde{B}_k$ .

Combining the above discussions, we get  $\bar{B}_k \subset \tilde{B}_k$ .

Conversely, assume that  $(\eta, \zeta) \in \tilde{B}_k$ . If  $(\eta, \zeta) \in \tilde{B}_{k-1} = \bar{B}_{k-1}$ , then it is immediately that  $(\eta, \zeta) \in \bar{B}_k$  by the definition of  $\bar{B}_n$ . Therefore, we may assume

$(\eta, \zeta) \notin \tilde{B}_{k-1}$ . Then, by (4.2), we can choose an

$$(\tilde{\eta}, \tilde{\zeta}) \in \bigcup_{l=0}^{k-1} \tilde{B}_l = \tilde{B}_{k-1} = \bar{B}_{k-1}$$

such that  $(\eta, \zeta) \rightarrow (\tilde{\eta}, \tilde{\zeta})$ . We say that  $(\tilde{\eta}, \tilde{\zeta}) \neq (\eta + e_u, \zeta)$ ,  $(\eta, \zeta + e_u)$  or  $(\eta + e_u, \zeta + e_u)$ . Otherwise, since  $(\eta, \zeta) = (\tilde{\eta} - e_u, \tilde{\zeta})$ ,  $(\tilde{\eta}, \tilde{\zeta} - e_u)$  or  $(\tilde{\eta} - e_u, \tilde{\zeta} - e_u)$  and  $(\tilde{\eta}, \tilde{\zeta}) \in \bar{B}_{k-1}$ , we would have  $(\eta, \zeta) \in \bar{B}_{k-2} \subset \bar{B}_{k-1} = \tilde{B}_{k-1}$ , this is a contradiction. So, by (4.1), there is a  $u \in S$  such that

$$(\eta, \zeta) = (\tilde{\eta} + e_u, \tilde{\zeta}), (\tilde{\eta}, \tilde{\zeta} + e_u) \text{ or } (\tilde{\eta} + e_u, \tilde{\zeta} + e_u).$$

By the definition of  $\bar{B}_{k-1}$ , we see that  $(\eta, \zeta) \in \bar{B}_k$ . Hence  $\tilde{B}_k \subset \bar{B}_k$ .

By induction, we have proved  $\tilde{B}_n = \bar{B}_n$  for each  $n \geq 0$ .

(iii) Suppose that  $(\eta, \zeta) \in \tilde{E}_n \subset \bar{B}_n$  and  $\tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) > 0$ . Then by (1.17)  $(\tilde{\eta}, \tilde{\zeta})$  should be one of  $(\eta \pm e_u, \zeta)$ ,  $(\eta, \zeta \pm e_u)$ ,  $(\eta \pm e_u, \zeta \pm e_u)$  ( $u \in S$ ),  $(\eta + e_u - e_v, \zeta)$ ,  $(\eta, \zeta + e_u - e_v)$  or  $(\eta + e_u - e_v, \zeta + e_u - e_v)$  ( $u, v \in S, u \neq v$ ). By (4.3),

$$(\tilde{\eta}, \tilde{\zeta}) \in \tilde{B}_{n+1} = \bigcup_{k=0}^{n+1} \tilde{E}_k,$$

and so (1.2) is satisfied.

The remains are obvious.

**Lemma 2.** Let  $\tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta}))$  be the  $Q$ -matrix corresponding to  $\tilde{Q}$  defined in (1.17). Then, for each  $(\eta, \zeta) \in \tilde{E}_n$  ( $n \geq 1$ ) there exist  $\{v_1, \dots, v_J\} \subset S$  and  $\{a_1, \dots, a_J\} \subset \mathcal{Z}_+$ ,  $\sum_{j=1}^J a_j = n$  such that

$$\sum_{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_{n-1}} \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) = \sum_{j=1}^J \delta(v_j, a_j). \tag{4.4}$$

Moreover,  $\sum_{j=1}^J a_j e_{v_j}$ , determined by the right side of (4.4) varies over the whole set  $\{\eta : |\eta| = n\}$  whenever  $(\eta, \zeta)$  varies over the whole set  $\tilde{E}_n$ .

*Proof* (i) By (4.3), we see that  $(\eta, \zeta) \in \tilde{E}_n = \tilde{B}_n \setminus \tilde{B}_{n-1}$  iff it has the form

$$\left( \sum_{i=1}^{k_i} e_{u_i}, \sum_{i=1}^i e_{u_i} + \sum_{l=k_i+1}^n e_l \right), \tag{4.5}$$

where  $i \in \{0, 1, \dots, n\}$ ,  $k_i \in \{i, i+1, \dots, n\}$ ,  $S \ni u_1, \dots, u_n$  may be repeated but

$$\{u_{i+1}, \dots, u_{k_i}\} \cap \{u_{k_i+1}, \dots, u_n\} = \phi.$$

(ii) Denote the distinct elements of  $\{u_1, \dots, u_i\} \setminus \{u_{i+1}, \dots, u_n\}$ ,  $\{u_{i+1}, \dots, u_{k_i}\}$  and  $\{u_{k_i+1}, \dots, u_n\}$  by  $\{v_1, \dots, v_{J_1}\}$ ,  $\{v_{J_1+1}, \dots, v_{J_2}\}$  and  $\{v_{J_2+1}, \dots, v_J\}$  respectively. Then we may write

$$\begin{cases} \sum_{i=1}^i e_{u_i} = \sum_{j=1}^{J_1} a_j e_{v_j} + \sum_{j=J_1+1}^J a_j^{(1)} e_{v_j}, & a_j > 0, a_j^{(1)} \geq 0, \\ \sum_{i=1}^{i+1} e_{u_i} = \sum_{j=J_1+1}^{J_2} a_j^{(2)} e_{v_j}, \\ \sum_{i=k_i+1}^n e_{u_i} = \sum_{j=J_2+1}^J a_j^{(2)} e_{v_j}, & a_j^{(2)} > 0. \end{cases} \tag{4.6}$$

Set  $a_j = a_j^{(1)} + a_j^{(2)}$ ,  $j = J_1+1, \dots, J$ . It follows from (i) and (4.6) that

$$\begin{cases} \eta = \sum_{j=1}^{J_1} a_j e_{v_j} + \sum_{j=J_1+1}^J a_j^{(1)} e_{v_j}, \\ \zeta = \sum_{j=1}^{J_1} a_j e_{v_j} + \sum_{j=J_1+1}^J a_j e_{v_j} + \sum_{j=J_1+1}^{J_2} a_j^{(1)} e_{v_j}, \end{cases} \quad (4.7)$$

and

$$\sum_{j=1}^J a_j = n. \quad (4.8)$$

(iii) We are going to compute  $\sum_{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_{n-1}} \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta}))$  for  $(\eta, \zeta) \in \tilde{E}_n$ . What

we need indeed is figure out

$$\tilde{E}_{(\eta, \zeta)} \equiv \{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_{n-1}: (\eta, \zeta) \rightarrow (\tilde{\eta}, \tilde{\zeta})\}.$$

By (4.1), (4.6), (4.7) and (i), we see that

(a)  $(\eta + e_{v_j}, \zeta + e_{v_j}), (\eta + e_{v_j}, \zeta), (\eta, \zeta + e_{v_j}) \notin \tilde{E}_{n-1}$ .

(b) If  $j \in \{1, \dots, J_1\}$ , then  $(\eta - e_{v_j}, \zeta - e_{v_j}) \in \tilde{E}_{n-1}$ ;  $(\eta - e_{v_j}, \zeta)$  and  $(\eta, \zeta - e_{v_j}) \in \tilde{E}_n$ .

(c) If  $j \in \{J_1+1, \dots, J_2\}$ , then  $(\eta - e_{v_j}, \zeta) \in \tilde{E}_{n-1}$ , either  $(\eta, \zeta - e_{v_j}) \in \tilde{E}_n$  (when  $a_j^{(1)} > 0$ ) or  $(\eta, \zeta - e_{v_j})$  has no meaning (when  $a_j^{(1)} = 0$ ), and either  $(\eta, \zeta) \rightarrow (\eta - e_{v_j}, \zeta - e_{v_j})$  (when  $a_j^{(1)} > 0$ ) or  $(\eta - e_{v_j}, \zeta - e_{v_j})$  has no meaning (when  $a_j^{(1)} = 0$ ).

(d) If  $j \in \{J_2+1, \dots, J\}$ , then  $(\eta, \zeta - e_{v_j}) \in \tilde{E}_{n-1}$ , either  $(\eta - e_{v_j}, \zeta) \in \tilde{E}_n$  (when  $a_j^{(1)} > 0$ ) or  $(\eta - e_{v_j}, \zeta)$  has no meaning (when  $a_j^{(1)} = 0$ ) and either  $(\eta, \zeta) \rightarrow (\eta - e_{v_j}, \zeta - e_{v_j})$  (when  $a_j^{(1)} > 0$ ) or  $(\eta - e_{v_j}, \zeta - e_{v_j})$  has no meaning (when  $a_j^{(1)} = 0$ ).

Combining the above discussion, we get

$$\begin{aligned} \sum_{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_{n-1}} \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) &= \sum_{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_{(\eta, \zeta)}} \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) \\ &= \sum_{j=1}^{J_1} \delta(v_j, \eta_{v_j}) \wedge \delta(v_j, \zeta_{v_j}) + \sum_{j=J_1+1}^{J_2} \delta(v_j, \eta_{v_j}) + \sum_{j=J_2+1}^J \delta(v_j, \zeta_{v_j}) \\ &= \sum_{j=1}^J \delta(v_j, a_j). \end{aligned}$$

(iv) The last assertion of Lemma 2 is obvious.

**Lemma 3.** Under the assumptions of Lemma 2, for each  $(\eta, \zeta) \in \tilde{E}_n$  ( $n \geq 0$ ), there exist  $\{v_1, \dots, v_J\} \subset S$  and  $\{a_1, \dots, a_J\} \subset Z_+$ ,  $\sum_{j=1}^J a_j = n$  such that

$$\sum_{j=1}^J \beta(v_j, a_j) \leq \sum_{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_{n+1}} \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) \leq 2 \left[ \sum_{j=1}^J \beta(v_j, a_j) + \sum_{u \neq v_j} \beta(u, 0) \right]. \quad (4.9)$$

Moreover,  $\sum_{j=1}^J a_j e_{v_j}$  varies over the whole  $\{\eta: |\eta| = n\}$  whenever  $(\eta, \zeta)$  varies over the whole  $\tilde{E}_n$ .

*Proof* Notice that (i) and (ii) in the proof of Lemma 2 are still available. Using the notations and the proof for (iii) given there, it follows that

$$\begin{aligned} \{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_{n+1}: (\eta, \zeta) \rightarrow (\tilde{\eta}, \tilde{\zeta})\} &= \{(\eta + e_u, \zeta + e_u): u \in \{v_{J_1+1}, \dots, v_J\}^c\} \\ &\cup \{(\eta + e_{v_j}, \zeta), (\eta, \zeta + e_{v_j}): j = J_1+1, \dots, J\}. \end{aligned}$$

By (1.17) and (4.7), we get

$$\begin{aligned} & \sum_{(\eta, \zeta) \in E_{n+1}} \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) \\ &= \sum_{\substack{u \neq v_j \\ j_1 < j_2}} \beta(u, \eta_u) \wedge \beta(u, \zeta_u) + \sum_{j=j_1+1}^J [\beta(v_j, \eta_{v_j}) + \beta(v_j, \zeta_{v_j})] \\ &= \sum_{j=1}^J \beta(v_j, a_j) + \sum_{j=j_1+1}^J \beta(v_j, a_j^{(1)}) + \sum_{\substack{u \neq v_j \\ 1 \leq j \leq J}} \beta(u, 0). \end{aligned}$$

Now (4.9) follows since  $\beta(u, k)$  is increasing in  $k$ .

**Lemma 4.** Under the assumptions of Lemma 2, for each  $n, k \in Z_+$  and each  $(\eta, \zeta) \in \tilde{E}_n$ , we have

$$\sum_{(\eta, \zeta) \in \tilde{E}_k} \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) = 0 \tag{4.11}$$

whenever  $|n - k| > 1$ .

*Proof* Using the proof of Lemma 2, it is easy to show that

$$\{(\tilde{\eta}, \tilde{\zeta}) \in \tilde{E}_k : (\eta, \zeta) \rightarrow (\tilde{\eta}, \tilde{\zeta})\} = \phi.$$

This proves (4.11).

*Proof of Theorem 5.* For the  $Q$ -matrix  $(q(\eta, \zeta) : \eta, \zeta \in X)$  defined by (1.16), take  $E_k = \{\eta \in X : |\eta| = k\}$ . Then the  $Q$ -matrix  $(q_{ij})$  defined by (1.1) is a birth-death  $Q$ -matrix and satisfies

$$\begin{cases} q_{k, k+1} = \max_{|\eta|=k+1} \{ \sum q(\eta, \zeta) : |\eta| = k \} = r_k, k \in Z_+, \\ q_{k, k-1} = \min_{|\eta|=k-1} \{ \sum q(\eta, \zeta) : |\eta| = k \} = s_k, k \in Z_+ \setminus \{0\}. \end{cases} \tag{4.12}$$

For the  $Q$ -matrix  $(\tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) : (\eta, \zeta), (\tilde{\eta}, \tilde{\zeta}) \in X \times X)$  replace  $E_k$  in (1.1) by  $\tilde{E}_k$  defined by (4.2), and define a  $Q$ -matrix  $(\tilde{q}_{ij})$  according to (1.1). By Lemma 4,  $(\tilde{q}_{ij})$  is again a birth-death  $Q$ -matrix. By Lemma 3 and Lemma 2, we have

$$\begin{cases} \tilde{r}_k \equiv \tilde{q}_{k, k+1} = \max_{(\eta, \zeta) \in \tilde{E}_{k+1}} \{ \sum \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) : (\eta, \zeta) \in \tilde{E}_k \} \\ \leq \max_{u \in S} \{ \sum 2\beta(u, \eta_u) : |\eta| = k \} = 2r_k \geq r_k, k \in Z_+, \\ \tilde{s}_k \equiv \tilde{q}_{k, k-1} = \min_{(\eta, \zeta) \in \tilde{E}_{k-1}} \{ \sum \tilde{q}((\eta, \zeta), (\tilde{\eta}, \tilde{\zeta})) : (\eta, \zeta) \in \tilde{E}_k \} \\ = \min_{u \in S} \{ \sum \delta(u, \eta_u) : |\eta| = k \} = s_k, k \in Z_+ \setminus \{0\}. \end{cases} \tag{4.13}$$

Hence, if (1.19) holds and  $r_k > 0$  ( $k \in Z_+$ ), then (4.12) and (4.13) imply that

$$\begin{aligned} \sum_{k=0}^{\infty} \left[ \frac{1}{r_k} + \frac{S_k}{r_k r_{k-1}} + \dots + \frac{S_k \dots S_1}{r_k \dots r_1 r_0} \right] &\geq \bar{R} = \infty, \\ \sum_{k=0}^{\infty} \left[ \frac{1}{\tilde{r}_k} + \frac{S_k}{\tilde{r}_k \tilde{r}_{k-1}} + \dots + \frac{S_k \dots S_1}{\tilde{r}_k \dots \tilde{r}_1 \tilde{r}_0} \right] &\geq \bar{R} = \infty. \end{aligned}$$

Therefore,  $\Omega$  and  $\tilde{\Omega}$  are regular by Theorem 1 and Theorem 3, (i). If (1.19) holds and  $r_0 = 0, r_k > 0$  ( $k \in Z_+ \setminus \{0\}$ ), then the conclusion follows from Theorem 1 and Theorem 3, (ii).

Finally, the remains are easily consequences of [10].

**Proposition 3.** Let  $S$  be finite. Then the  $Q$ -matrices defined by (1.12), (1.13)

and (1.14) satisfy the assumptions of Theorem 5.

*Proof* The calculations are elementary once we write down the Q-matrices.

For (1.12)

$$\begin{cases} \beta(u, \eta_u) = \lambda_1 a_u \begin{pmatrix} \eta_u \\ 2 \end{pmatrix} + \lambda_4 b_u, \delta(u, \eta_u) = \lambda_2 \begin{pmatrix} \eta_u \\ 3 \end{pmatrix} + \lambda_3 \eta_u, \\ \gamma(u, v, \eta_u, \eta_v) = \eta_u p(u, v), u, v \in S, u \neq v. \end{cases} \quad (4.14)$$

For (1.13) and (1.14),  $S$  and  $u$  should be replaced with  $S \times \{1, 2\}$  and  $(u, i)$  ( $u \in S, i=1, 2$ ) respectively. Then, for (1.13), we have

$$\begin{aligned} \beta((u, i), \eta_{ui}) &= \begin{cases} \lambda_1 a_u \eta_{u1}, i=1, \\ 0, i=2, \end{cases} \\ \delta((u, i), \eta_{ui}) &= \begin{cases} 0, i=1, \\ \lambda_3 b_u \eta_{u2}, i=2, \end{cases} \\ \gamma((u, i), (v, j), \eta_{ui}, \eta_{vj}) &= \begin{cases} \lambda_2 \eta_{u1} \eta_{v2}, u=v, i=1, j=2, \\ \eta_{u1} p_1(u, v), u \neq v, i=j=1, \\ \eta_{u2} p_2(u, v), u \neq v, i=j=2, \\ 0, \text{other } (u, i) \neq (v, j), \end{cases} \end{aligned}$$

and for (1.14), we have

$$\begin{aligned} \beta((u, i), \eta_{ui}) &= \begin{cases} \lambda_1 a_u, i=1, \\ 0, i=2, \end{cases} \\ \delta((u, i), \eta_{ui}) &= \begin{cases} \lambda_4 \eta_{u1}, i=1, \\ 0, i=2, \end{cases} \\ \gamma((u, i), (v, j), \eta_{ui}, \eta_{vj}) &= \begin{cases} \lambda_2 b_u \eta_{u1}, u=v, i=1, j=2, \\ \lambda_3 \begin{pmatrix} \eta_{u1} \\ 2 \end{pmatrix} \eta_{u2}, u=v, i=2, j=1, \\ \eta_{ui} p_i(u, v), u \neq v, i=j=1, 2, \\ 0, \text{other } (u, i) \neq (v, j). \end{cases} \end{aligned}$$

## § 5. Recurrence

**Lemma 1.** Let  $(q_{ij}; i, j \in E)$  be a regular irreducible Q-matrix. Then, the  $(q_{ij})$ -process is recurrent iff

$$x_i = \sum_{k \neq j_0} \bar{p}_{ik} x_k, \quad 0 \leq x_i \leq 1, \quad i \in E \quad (5.1)$$

has only zero solution for some  $j_0 \in E$  (or equivalently, for any  $j_0 \in E$ ), where  $(p_{ij})$  is the jump matrix of  $(q_{ij})$ , i.e.,

$$\bar{p}_{ij} = \begin{cases} q_{ij}/q_i, & i \neq j, \\ 0, & i=j, i, j \in E. \end{cases} \quad (5.2)$$

*Proof* Clearly, we can fix a  $j_0 \in E$ . By [7, Theorem 1], the  $(q_{ij})$ -process is recurrent iff the jump chain  $(\bar{p}_{ij})$  is recurrent. Then, by [4, Theorem 6.6.1] it is equivalent to that the minimal nonnegative solution of



$$x_i = \sum_{k \neq j_0} \bar{p}_{ik} x_k + \bar{p}_{i0}, \quad i \in E, \quad (5.3)$$

is  $x_i^* = 1$  ( $i \in E$ ). Therefore, by [4, Theorem 5.6.3], the proof is reduced to showing that (5.3) has no nonnegative non-constant bounded solution iff (5.1) has only zero solution. To this end, let  $(\tilde{x}_i: i \in E)$  be a non-zero solution of (5.1). Then  $(y_i = 1 + \tilde{x}_i: i \in E)$  is a nonnegative non-constant bounded solution of (5.3). Conversely, if (5.3) has such a solution, then the minimal nonnegative solution of (5.3) satisfies  $x_i^* \leq 1$  and there exists at least an  $i_0 \in E$  so that  $x_{i_0}^* < 1$ , since  $(y_i = 1, i \in E)$  is a solution of (5.3). Hence,  $(\tilde{x}_i = 1 - x_i^*: i \in E)$  is a non-zero solution of (5.1).

*Proof of Theorem 6.* By Lemma 1, to prove the first part of Theorem 6, it suffices to show that (5.1) has only zero solution implies that

$$u(\eta) = \sum_{\zeta \neq \theta} \bar{p}(\eta, \zeta) u(\zeta), \quad 0 \leq u(\eta) \leq 1, \quad \eta \in X_0 \quad (5.4)$$

has only zero solution, where  $(\bar{p}(\eta, \zeta))$  is the jump matrix of  $(q(\eta, \zeta))$ . Now, the proof is similar to the proof of Theorem 1. Suppose that  $\{u(\eta): \eta \in X_0\}$  is a non-zero solution to (5.4). Then

$$q(\eta)u(\eta) = \sum_{\zeta \neq \theta, \eta} q(\eta, \zeta)u(\zeta), \quad 0 \leq u(\eta) \leq 1, \quad \eta \in X_0. \quad (5.5)$$

Set  $u_k = \sup\{u(\eta): \eta \in E_k\}$ ,  $i \in Z_+$ . Since  $E_0 = \{\theta\}$ ,  $E_k$  is a finite set ( $k \geq 1$ ), there exists an  $\eta^{(k)} \in E_k$  ( $k \geq 1$ ) so that

$$u_0 = u(\theta), \quad u_k = u(\eta^{(k)}), \quad (5.6)$$

$$q_0 u_0 = q(\theta)u(\theta) = \sum_{\zeta \neq \theta} q(\theta, \zeta)u(\zeta) \leq \sum_{\zeta \in E_1} q(\theta, \zeta)u_1 = q_{01}u_1. \quad (5.7)$$

If  $k \geq 1$ , by using the method in the proof of (2.6), it follows that

$$q(\eta^{(k)}, \theta)u_k + \sum_{j=1}^{k-1} \left( \sum_{\zeta \in E_j} q(\eta^{(k)}, \zeta) \right) (u_k - u_j) \leq \sum_{\zeta \in E_{k+1}} q(\eta^{(k)}, \zeta) (u_{k+1} - u_k),$$

so  $u_k \uparrow$ , and

$$q_{k0}u_k + \sum_{j=1}^{k-1} q_{kj} (u_k - u_j) \leq q_{k, k+1} (u_{k+1} - u_k), \quad k \geq 1. \quad (5.8)$$

Combining (5.7) and (5.8), we get

$$q_k u_k \leq \sum_{j \neq 0, k} q_{kj} u_j, \quad k \in Z_+.$$

It is now easy to show that  $u_i = 0$ ,  $i \in Z_+$  (cf. the proof of § 2, Lemma 1) since (5.1) has only zero solution. But this is a contradiction to  $u(\eta) \neq 0$ .

As for the second part of Theorem 6, by Lemma 1, we need only to prove that

$$x_i = \sum_{k \neq 0, i} \frac{q_{ik}}{q_i} x_k, \quad 0 \leq x_i \leq 1, \quad i \in Z_+ \quad (5.9)$$

has non-zero solution iff  $\sum_{k=0}^{\infty} F_k^{(0)} < \infty$ . Notice that (5.9) has non-zero solution iff

$$x_i = \sum_{k \neq 0, i} \frac{q_{ik}}{q_i} x_k, \quad 0 \leq x_i \leq 1, \quad i \in Z_+ \quad (5.10)$$

has nonnegative bounded solution, and (5.10) has at most one solution which is

$$\tilde{x}_0 = 1 = \tilde{x}_1, \quad (5.11)$$

$$\tilde{x}_{i+1} - \tilde{x}_i = \left( \sum_{j=1}^{i-1} q_{ij} (\tilde{x}_i - \tilde{x}_j) + q_{i0} \tilde{x}_i \right) / q_{i,i+1}, \quad i \geq 1.$$

Clearly,  $(\tilde{x}_i)$  is increasing. Thus, the proof is reduced to showing that  $(\tilde{x}_i)$  is bounded iff

$$\sum_{k=0}^{\infty} F_k^{(0)} < \infty.$$

Observe that

$$\sum_{j=1}^{i-1} q_{ij}^{(j)} (\tilde{x}_{i+1} - \tilde{x}_j) = \sum_{j=1}^{i-1} q_{ij} (\tilde{x}_i - \tilde{x}_j).$$

Hence

$$\tilde{x}_{i+1} - \tilde{x}_i = \sum_{j=1}^{i-1} \frac{q_j^{(j)}}{q_{i,i+1}} (\tilde{x}_{i+1} - \tilde{x}_j) + \frac{q_i^{(0)}}{q_{i,i+1}}, \quad i \geq 1.$$

This proves that  $(y_i \equiv \tilde{x}_{i+1} - \tilde{x}_i; i \geq 1)$  is a nonnegative finite solution to

$$z_i = \sum_{j=1}^{i-1} \frac{q_j^{(j)}}{q_{i,i+1}} z_j + \frac{q_i^{(0)}}{q_{i,i+1}}, \quad i \geq 1. \quad (5.12)$$

On the other hand, the solution to (5.12) is unique, i.e.,

$$z_1^* = q_1^{(0)} / q_{12},$$

$$z_i^* = \sum_{j=1}^{i-1} (q_j^{(j)} / q_{j,j+1}) z_j^* + q_i^{(0)} / q_{i,i+1}, \quad i \geq 1,$$

and it is easy to show

$$z_i^* = F_i^{(0)}, \quad i \geq 1$$

by induction. Using the above remarks, we get

$$\tilde{x}_0 = \tilde{x}_1 = 1 = F_0^{(0)}, \quad \tilde{x}_{i+1} - \tilde{x}_i = F_i^{(0)}, \quad i \geq 1.$$

This of course implies what we need.

**Proposition 4.** *Let  $S$  be finite. Then the  $(q(\eta, \zeta))$ -process defined by (1.12) is recurrent.*

*Proof* Take  $E_k = \{\eta \in X: |\eta| = k\}$ . By (3.4) and (3.6), we have  $s_{k+1}/r_{k+1} \rightarrow \infty$  ( $k \rightarrow \infty$ ). Hence

$$\sum_{k=0}^{\infty} F_k^{(0)} = 1 + \sum_{k=1}^{\infty} \frac{s_k \cdots s_1}{r_k \cdots r_1} = \infty.$$

Now Theorem 6 is available.

## § 6. Positive Recurrence and Ergodicity

*Proof of Theorem 7.* If the  $(q_{ij})$ -process is positive recurrent, then for some (each)  $i_0 \in Z_+$  there exists a nonnegative solution to

$$x_i \geq \sum_{j=i_0, i} \frac{q_{ij}}{q_i} x_j + \frac{1}{q_i}, \quad i \neq i_0, \quad (6.1)$$

$$\sum_{j \neq i_0} \frac{q_{ij}}{q_{i_0}} x_j < \infty$$

by [4, Theorem 9.4.1]. But (6.1) is just the condition (1.22) if we set  $x_{i_0} = 0$ , in addition. Therefore (1.22) is necessary.

Conversely, if  $\{u_i: i \in Z_+\}$  is a nonnegative solution of (1.22) for an  $i_0 \in Z_+$ , we define

$$c_i = 1, i \neq i_0; = - \sum_j q_{ij} u_j, i = i_0. \tag{6.2}$$

Then  $c_i + \sum_j q_{ij} u_j \leq 0, i \in Z_+$ , and so

$$u_i - c \geq \sum_{j \neq i} \frac{q_{ij}}{\lambda + q_{ij}} (u_j - c) + \frac{c_i - \lambda c}{\lambda + q_{ij}}, i \in Z_+, \tag{6.3}$$

where  $c = (\inf\{c_j/\lambda: j \in Z_+\}) \wedge 0 > -\infty$ . Denote the Laplace transform of the minimal  $(q_{ij})$ -process by  $(P_{ij}^{\min}(\lambda))$ . It follows from [4, Theorem 3.3.3 and Theorem 3.3.1] that

$$u_i - c \geq \sum_j P_{ij}^{\min}(\lambda) (c_j - \lambda c). \tag{6.4}$$

Since  $(q_{ij})$  is regular,  $\lambda \sum_j P_{ij}^{\min}(\lambda) = 1, i \in Z_+, \lambda > 0$ , it follows from (6.2) and (6.4) that

$$u_i \geq \sum_j P_{ij}^{\min}(\lambda) c_j = \sum_{i \neq i_0} P_{ij}^{\min}(\lambda) + P_{i_0 i}^{\min}(\lambda) c_{i_0} = \lambda^{-1} + P_{i_0 i}^{\min}(\lambda) (c_{i_0} - 1),$$

i.e.,

$$\lambda u_i \geq 1 + \lambda P_{i_0 i}^{\min}(\lambda) (c_{i_0} - 1), i \in Z_+.$$

Because of  $\lim_{\lambda \rightarrow 0} \lambda P_{ij}^{\min}(\lambda) = \lim_{t \rightarrow \infty} p_{ij}(t) = \pi_{ij}, i, j \in Z_+$ , we have

$$0 \geq 1 + \pi_{i_0 i} (c_{i_0} - 1), i \in Z_+,$$

and so  $\pi_{i_0 i} \neq 0$  for each  $i \in Z_+$ . This implies that  $\pi_i = \pi_{ij} > 0, i, j \in Z_+$  and  $\sum_i \pi_i = 1$ . So the  $(q_{ij})$ -process is ergodic.

*Proof of Theorem 8* By Theorem 7, it suffices to prove that

$$\begin{aligned} \sum_{\zeta} q(\eta, \zeta) u(\zeta) + 1 &\leq 0, \eta \neq \theta, \\ |\sum_{\zeta} q(\theta, \zeta) u(\zeta)| &< \infty \end{aligned} \tag{6.5}$$

has nonnegative solution. To this end, put

$$u(\eta) = u_k, \eta \in E_k, k \in Z_+.$$

Given  $\eta \notin E_0$ , there exists only one  $k \in Z_+ \setminus \{0\}$  so that  $\eta \in E_k$ . Hence

$$\begin{aligned} \sum_{\zeta} q(\eta, \zeta) u(\zeta) + 1 &= - \sum_{j=0}^{k-1} \sum_{\zeta \in E_j} q(\eta, \zeta) (u_k - u_j) + \sum_{\zeta \in E_{k+1}} q(\eta, \zeta) (u_{k+1} - u_k) + 1 \\ &< - \sum_{j=0}^{k-1} q_{kj} (u_k - u_j) + q_{k, k+1} (u_{k+1} - u_k) + 1 = \sum_j q_{kj} u_j + 1 \leq 0 \end{aligned}$$

by (1.1), (1.2),  $u_k \nearrow$  and the assumption of this theorem. But

$$\sum_{\zeta} q(\theta, \zeta) u(\zeta) = \sum_{\zeta \in E_1} q(\theta, \zeta) u_1 = q_{01} u_1$$

is finite, so (6.5) holds.

*Proof of Theorem 9* Let  $\{u_i: i \in Z_+\}$  be a nonnegative increasing solution to (1.22) with  $i_0=0$ . Then

$$q_{01}(u_1 - u_0) = \left| \sum_j q_{0j} u_j \right| < \infty, \tag{6.6}$$

$$\sum_j q_{kj} u_j + 1 \leq 0, \quad k > 0. \tag{6.7}$$

By (1.7), (6.7) and Abelian transform, we get

$$q_{k, k+1}(u_{k+1} - u_k) + 1 \leq \sum_{j=0}^{k-1} q_{kj}(u_k \cdots u_j) = \sum_{j=0}^{k-1} q_k^{(j)}(u_{j+1} - u_j). \tag{6.8}$$

Put  $v_k = u_{k+1} - u_k (k \geq 0)$ . Then  $v_0$  is finite by (6.6). By (1.9), (1.25), (6.8) and induction it follows that

$$v_k \leq F_k^{(0)} v_0 - d_k, \quad k \in Z_+. \tag{6.9}$$

Now, (1.24) follows from (6.9) since  $u_k \uparrow, v_k \geq 0, k \in Z_+$ .

Conversely, if (1.24) holds, then (6.6) holds for  $(u_k: k \in Z_+)$  defined by (1.26). Hence, by (1.9) and (1.25), for each  $k > 0$ , we have

$$\begin{aligned} 1 + q_{k, k+1}(u_{k+1} - u_k) &= q_{k, k+1} F_k^{(0)} u_1 - q_{k, k+1} d_k + 1 = \sum_{s=0}^{k-1} q_k^{(s)} F_s^{(0)} u_1 - \sum_{s=0}^{k-1} q_k^{(s)} d_s \\ &= \sum_{s=0}^{k-1} q_k^{(s)} (F_s^{(0)} u_1 - d_s) = \sum_{s=0}^{k-1} q_k^{(s)} (u_{s+1} - u_s) = \sum_{j=0}^{k-1} q_{kj}(u_k - u_j), \end{aligned}$$

i. e.  $\sum_j q_{kj} u_j + 1 = 0$ , and so (1.22) has a nonnegative increasing solution when  $i_0 = 0$ .

**Corollary.** Let  $(q_{ij})$  be a birth-death Q-matrix:

$$r_k = q_{k, k+1} > 0, \quad k \geq 0,$$

$$s_k = q_{k, k-1} > 0, \quad k \geq 1.$$

Then, the condition (1.24) is equivalent to

$$\sum_{k=0}^{\infty} \frac{r_0 \cdots r_k}{s_1 \cdots s_{k+1}} < \infty. \tag{6.11}$$

*Proof* By (1.9) and (1.25), we have

$$\frac{d_k}{F_k^{(0)}} = \frac{r_1 \cdots r_k}{s_1 \cdots s_k} \cdot \frac{1}{r_k} [1 + s_k d_{k-1}] = \cdots = \frac{1}{r_0} \sum_{l=0}^{k-1} \frac{r_0 r_1 \cdots r_l}{s_1 s_2 \cdots s_{l+1}}.$$

**Proposition 5.** Let  $S$  be finite. Then the Q-process corresponding to Schlögl model is ergodic.

*Proof* By (3.4) and (3.6),  $\frac{r_k}{s_k} \rightarrow 0 (k \rightarrow \infty)$ . Hence (1.24) holds.

*Proof of Theorem 10* Let  $(u_i: i \in Z_+)$  be a nonnegative solution to (1.22) with  $i_0=0$ , and put  $v_k = u_{k+1} - u_k$ . Then (6.9) holds, and so

$$0 \leq u_{k+1} = \sum_{i=0}^k v_i + u_0 \leq \left( \sum_{i=0}^k F_i^{(0)} \right) v_0 - \sum_{i=0}^k d_i + u_0 \leq \left( \sum_{i=0}^k F_i^{(0)} \right) u_1 - \sum_{i=0}^k d_i.$$

Therefore, we have  $\hat{d} \leq u_1 < \infty$ . Conversely, if (1.27) holds, we choose  $u_0 = 0, u_1 \geq \hat{d}$ , and define  $u_k (k \geq 2)$  according to (1.26). Then, by using the proof of Theorem 9, it follows that

$$\sum_j q_{ij}u_j + 1 = 0, \quad k \neq 0,$$

$$\sum_j q_{0j}u_j = q_{01}u_1 \text{ is finite,}$$

and

$$u_k = \sum_{l=0}^{k-1} (F_l^{(0)}u_1 - d_l) \geq d \sum_{l=0}^{k-1} F_l^{(0)} - \sum_{l=0}^{k-1} d_l \geq 0, \quad k \geq 2.$$

We get a nonnegative solution to (1.22) with  $i_0 = 0$ .

The last assertion of Theorem 10 is obvious.

**Acknowledgment.** We would like to thank those people who attended our seminar for their discussion with us, in particular, Ms. Lu Quan, for her checking some examples in this paper.

### References

- [1] Nicolis, G. and Prigogine, I., Self-organization in Non-equilibrium Systems, *John Wiley & Sons*, 1977.
- [2] Haken, H., *Synergetics*, Springer-Verlag, 1977.
- [3] 严士健、李占柄, 非平衡系统的概率模型及 Master 方程的建立. *物理学报*, 29 (1980), 139—152
- [4] 侯振挺、郭青峰, 齐次可列马尔可夫过程, 科学出版社, 北京, 1978.
- [5] Reuter, G. E. H., Competition processes, *Proc. of the 4th Berkeley Symposium on Math. Statis. and Prob.* 1961. Vol. 2. 421—430.
- [6] 张建康, 广生灭过程的性质, *数学物理学报*, 4:2(1984), 241—259.
- [7] 吴立德, 可数马尔科夫过程的状态的分类, *数学学报*, 15 (1965), 32—41
- [8] 王梓坤, 生灭过程与马尔科夫链, 科学出版社, 北京, 1980.
- [9] 侯振挺,  $Q$ -过程唯一性准则, 湖南科学技术出版社, 长沙 1982.
- [10] 陈木法, 马链的基本耦合, *北京师范大学学报*, 4(1984), 3—10.