

Algebraic structure of the Gaussian-PDMF space and applications on fuzzy equations

Chuang Zheng*

January 23, 2024

Abstract

In this paper, we extend the research presented in [26] by establishing the algebraic structure of the Gaussian Probability Density Membership Function (Gaussian-PDMF) space. We consider fixed objective and subjective entities, denoted as (h, p) , and provide the explicit form of the membership function. Consequently, every fuzzy number with the membership function in $X_{h,p}(\mathbb{R})$, denoted as \tilde{x} , can be uniquely identified by a vector $\langle x; d^-, d^+, \mu^-, \mu^+ \rangle$. Here, $x \in \mathbb{R}$ represents the “leading factor” of the fuzzy number \tilde{x} with a membership degree equal to 1. The parameters d^- (left side) and d^+ (right side) denote the lengths of the compact support, while μ^- (left side) and μ^+ (right side) represent the shapes. We introduce five operators: addition, subtraction, multiplication, scalar multiplication, and division. We demonstrate that, based on our definitions, the Gaussian-PDMF space exhibits a well-defined algebraic structure. For instance, $X_{h,p}(\mathbb{R})$ is a vector space over \mathbb{R} , featuring a subspace that forms a division ring, allowing for the representation of fuzzy polynomials, among other properties. We provide several examples to illustrate our theoretical results.

Key Words. Fuzzy numbers; Gaussian probability density membership function (Gaussian-PDMF) space; Fuzzy arithmetic; Abelian group; Ring; Fuzzy Vieta’s formula; Linear space; Fuzzy equations.

1 Introduction

Fuzzy numbers and fuzzy set theory are topics originated from Zadeh ([33]) by dealing with the imprecise quantities and uncertainty. Since then, they have found successful applications

*School of Mathematical Sciences, Beijing Normal University, 100875 Beijing, China. Email: chuang.zheng@bnu.edu.cn.

in a wide range of areas, including pure and applied mathematics, computer science, and related fields such as fuzzy logic, fuzzy information, soft computing, and fuzzy control.

In general, a fuzzy number can be uniquely determined by its membership function. The construction of an appropriate membership function is the cornerstone upon which fuzzy set theory has evolved. For existing references, see [6, 16, 19, 27, 29, 30] and the references therein. Particularly, in the realm of artificial intelligence, linguistic variables are associated with a membership function based on subjective choices (see, for instance, [3, 17]).

Once membership functions of fuzzy numbers are identified, a fundamental concern is performing arithmetic operations on them. In the fuzzy world, arithmetic operations on real numbers at the classical crisp set level translate into algebraic operations on membership functions. Numerous papers have explored the fundamental definitions of these operations and the corresponding algebraic structures. Triangular norms, for example, address binary operations on the interval $[0, 1]$ ([15]). Interactive fuzzy numbers and their arithmetic operators are proposed through joint possibility distributions ([32, 8, 10]), with a notable method offering minimum norm via the sup-J extension principle ([11]). Intuitionistic fuzzy sets, along with probabilistic addition, have successfully addressed fuzzy aggregation problems in expert systems ([1, 2, 24, 31]). Moreover, in the context of fuzzy differential equations, a common definition of (generalized) Hukuhara differentiability necessitates designing the difference between two fuzzy numbers using α -cuts of their corresponding membership functions ([9, 14, 20]). Applications encompass various classes of membership functions such as triangle MF, trapezoidal MF, Sigmoid MF, among others. MATLAB's standard toolbox offers more than 11 different types of membership functions for constructing fuzzy models.

In [26], a new class of membership functions is introduced through the composition of two types of nonlinear functions: $h : [0, 1] \rightarrow \mathbb{R}$ and $p : \mathbb{R} \rightarrow [0, 1]$. Here, h represents subjective perception, while p provides objective information via the probability density function. This class of functions defines the Probability Density Membership Function (PDMF) space $X_{h,p}(\mathbb{R})$. Specifically, h is chosen as the tangent function, and p is given by the Gaussian kernel $p(\cdot, \mu)$ with $\sigma = 1$ and μ undetermined, where μ measures fuzziness, and arithmetic operations are designed via μ to achieve a linear structure. Such class of MFs is called as the Gaussian-PDMFs. Notably, the α -cuts representation is unnecessary during computation implementation.

In this paper, we introduce a new notation of the fuzzy number \tilde{x} with the membership function in Gaussian-PDMF space $X_{h,p}(\mathbb{R})$. As we show later, \tilde{x} is uniquely identified by a vector $[x; d^-, d^+, \mu^-, \mu^+]$, where $x \in \mathbb{R}$ represents the "leading factor" of the fuzzy number \tilde{x} with membership degree equaling to 1, d^- (resp. d^+) is the length of the compact support on the left side (resp. right side) and μ^- (resp. μ^+) represents the shape of the left (resp. right) side. We design five algebraic operations: addition, subtraction, multiplication, division and scalar multiplication in terms of the five parameters of \tilde{x} .

The first result is that addition \oplus forms an abelian group on $X_{h,p}(\mathbb{R})$. Due to the existence of the inverse element, the minus \ominus can be seen, or directly defined by, the addition of the inverse element. Furthermore, together with the multiplication \otimes , $X_{h,p}(\mathbb{R})$ turns to a

commutative ring with identity. Following the standard procedure of the algebra, we state several basic properties of the space $X_{h,p}(\mathbb{R})$ with two operations \oplus and \otimes . Collecting those elements with inverse under multiplication, we form a subset of $X_{h,p}(\mathbb{R})$, denoted by $U(X)$. As a result, the cancellation law holds with respect to \oplus and \otimes . We give several applications for this algebraic structure, such as the solvability of fuzzy equations, Binomial Theorem, fuzzy Vieta's formula, etc. The last scalar multiplication on $X_{h,p}(\mathbb{R})$ is the basic operation for the linear algebra. By constructing a basis of $X_{h,p}(\mathbb{R})$, we give the coordinates of \tilde{x} and shows that the coordinate mapping is a isomorphism from $X_{h,p}(\mathbb{R})$ onto \mathbb{R}^5 .

The structure of the paper is organized as follows. In Section 2, we present the basic concepts of fuzzy numbers and outline the requirements of the membership function. In Section 3, we provide exact definitions of arithmetic operators on $X_{h,p}(\mathbb{R})$, along with the algebraic structure such as abelian group, commutative ring with identity, etc. Additionally, we introduce a binomial theorem and demonstrate its application to second-order fuzzy equations. In Section 4, we prove that $X_{h,p}(\mathbb{R})$ is a vector space and establish the isomorphism between $X_{h,p}(\mathbb{R})$ and \mathbb{R}^5 . In Section 5, we present numerical examples and corresponding graphs to illustrate the operations under consideration. Finally, in Section 6, we offer a concluding remark to provide a comprehensive summary of the paper.

2 Preliminary

The formulation of membership functions is the crucial step in the design of fuzzy system. There are several methods to develop them. We summarize some of them as follows:

- 1) L-R linear functions, which is the simplest possible model ([7]);
- 2) Rational functions of polynomials ([12, 13]);
- 3) B-Spline MF ([25]);
- 4) Piecewise linear functions ([28] and refs [8-20] in it);
- 5) Gaussian PDMF and the corresponding Gaussian-PDMF space $X_{h,p}(\mathbb{R})$ ([26]).

In all of these definitions of fuzzy numbers, the membership function needs to satisfy the following assumptions:

Definition 2.1 *A fuzzy number A is a fuzzy subset of the real line \mathbb{R} with membership function f_A which possesses the following properties:*

- a) f_A is fuzzy convex,
- b) f_A is normal i.e., $\exists x_0 \in \mathbb{R}$ such that $f_A(x_0) = 1$,
- c) f_A is upper semi-continuous,

d) The closure of the set $\{x \in \mathbb{R} | f_A(x) > 0\}$ is compact.

Definition 2.1 is straightforward and has been used extensively in practical applications ([22]). However, the above conditions are too vague and a particular class of functions, named as monotonic fuzzy numbers, is introduced with the following more precise assumptions:

Definition 2.2 A monotonic fuzzy number \tilde{b} , denoted by $\tilde{b} = (a, b, c)$, is defined as a membership function $f(x)$ which possesses the following properties ([7]):

- a) $f(x)$ is increasing on the interval $[a, b]$ and decreasing on $[b, c]$,
- b) $f(x) = 1$ for $x = b$, $f(x) = 0$ for $x \leq a$ or $x \geq c$,
- c) $f(x)$ is upper semi-continuous,

where a, b, c , are real numbers satisfying $-\infty < a < b < c < +\infty$.

Clearly, a class of triangular fuzzy numbers is a subset of the class of monotonic fuzzy numbers. It is due to the fact that, in the definition of the triangular fuzzy numbers, the function in the condition a) of Definition 2.2 is restricted by linear ones ([18]).

Recently, a new class of fuzzy numbers called PDMF is given in [26], in which extra information between a, b and b, c are clarified. More precisely, a monotonic number \tilde{b} , denoted by $\tilde{b} = \langle (a, b, c); P(x^-, y^-), Q(x^+, y^+) \rangle$, is defined as a membership function $f(x)$ which possesses the following structure:

$$f(x) = \begin{cases} 0, & x \in (-\infty, a] \\ \int_{-\infty}^{h^-(x)} p^-(y) dy, & x \in (a, b) \\ 1, & x = b \\ \int_{-\infty}^{h^+(x)} p^+(y) dy, & x \in (b, c) \\ 0, & x \in [c, +\infty). \end{cases} \quad (2.1)$$

In (2.1), h^-, h^+ are the auxiliary functions satisfying

i) $\lim_{x \rightarrow a^+} h^-(x) = \lim_{x \rightarrow c^-} h^+(x) = -\infty$, $\lim_{x \rightarrow b^-} h^-(x) = \lim_{x \rightarrow b^+} h^+(x) = +\infty$,

ii) h^- is continuous and increasing on (a, b) , h^+ is continuous and decreasing on (b, c) ,

and p^-, p^+ are probability density functions (PDFs) satisfying

$$\int_{-\infty}^{+\infty} p^-(y) dy = 1, \int_{-\infty}^{+\infty} p^+(y) dy = 1, p^-(t) \geq 0, p^+(t) \geq 0, \quad \forall t \in (-\infty, +\infty).$$

· Moreover, it is required that the shape of $f(x)$ passes through P, Q .

The following Theorem stated in [26] provides the well-posedness of the definition (2.1) for $f(x)$:

Theorem 2.1 *Let*

$$h^-(x) = h\left(\frac{x-a}{b-a}\right), h^+(x) = h\left(\frac{c-x}{c-b}\right) \quad \text{with} \quad \lim_{x \rightarrow 0^+} h(x) = -\infty, \lim_{x \rightarrow 1^-} h(x) = +\infty,$$

and $h(x)$ is continuous and increasing on $(0, 1)$. Let p^- and p^+ be originated from the same class p of PDFs. Then there exists at least one pair (h, p) such that the graph of $f_{h,p}$ passes through P and Q , i.e., $f_{h,p}(x^-) = y^-$ and $f_{h,p}(x^+) = y^+$. Moreover, the PDMF in the space

$$X_{h,p}(\mathbb{R}) := \{f_{h,p}(x) : \mathbb{R} \rightarrow [0, 1] \text{ is as the form of (2.1) : } a < b < c\}$$

fulfills all requirements in Definition 2.1 and Definition 2.2.

In this paper, to specify the class of functions under consideration, we fix h as the tangent function and p is given by the Gaussian kernel $p(\cdot, \mu)$ with $\sigma = 1$ and μ to be undetermined, i.e.

$$h(x) = \tan\left(\pi x - \frac{\pi}{2}\right), \quad x \in (0, 1), \quad p(t) = p(t; \mu) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(t-\mu)^2}, \quad t \in \mathbb{R}. \quad (2.2)$$

The two control points $P(x^-, y^-), Q(x^+, y^+)$ are placed on each side of the central point $B(x_0, 1)$, and the shape of the function passes through $A(x_0 - d^-, 0)$ and $C(x_0 + d^+, 0)$ with $d^- = x_0 - a$ and $d^+ = c - x_0$, respectively. See Figure 2 for the sample of the shape of the membership function. Note that here d^- and d^+ are the deviation measures from the center x_0 to the left side and right side, respectively. Clearly, taking into account (2.2), the function in the PDMF space has the explicit form

$$f_{h,p}(x) = \begin{cases} 0, & x \in (-\infty, x_0 - d^-] \\ f_-(x; x_0, d^-, \mu^-), & x \in (x_0 - d^-, x_0) \\ 1, & x = x_0 \\ f_+(x; x_0, d^+, \mu^+), & x \in (x_0, x_0 + d^+) \\ 0, & x \in [x_0 + d^+, +\infty) \end{cases} \quad (2.3)$$

where f_- and f_+ are given by

$$\begin{aligned} f_-(x; x_0, d^-, \mu^-) &= \int_{-\infty}^{\tan\left(\frac{\pi}{d^-}(x-x_0+d^-)-\frac{\pi}{2}\right)} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(t-\mu^-)^2} dt, & x \in (x_0 - d^-, x_0), \\ f_+(x; x_0, d^+, \mu^+) &= \int_{-\infty}^{\tan\left(\frac{\pi}{d^+}(x_0+d^+-x)-\frac{\pi}{2}\right)} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(t-\mu^+)^2} dt, & x \in (x_0, x_0 + d^+). \end{aligned} \quad (2.4)$$

The function as above is the **Gaussian-PDMF** and the corresponding function space is the **Gaussian-PDMF Space**. More precisely,

$$X_{h,p}(\mathbb{R}) := \{f_{\tan,G}(x) : \mathbb{R} \rightarrow [0, 1] \text{ is as the form of (2.3) with (2.4)}\}. \quad (2.5)$$

As described in Theorem 4.1 of [26], there is a unique vector $\langle(x_0 - d^-, x_0, x_0 + d^+); \mu^-, \mu^+\rangle$ for each membership function $f_{\text{tan},G}(x)$ in the Gaussian-PDMF space passing through the two control points $P(x^-, y^-), Q(x^+, y^+)$.

Consequently, it is reasonable to give two equivalent notations of the Gaussian-PDMF as

$$\langle(x_0 - d^-, x_0, x_0 + d^+); P, Q\rangle \iff \langle x_0; d^-, d^+, \mu^-, \mu^+\rangle. \quad (2.6)$$

Here, (μ^-, μ^+) can be uniquely identified by means of the inverse functions of Formula (2.4)




Figure 1: Fuzzy number $\tilde{1} = \langle(-1, 1, 2); P, Q\rangle$ or $\langle 2; 2, 1, \mu^-, \mu^+\rangle$ where μ^- is uniquely identified by the point P via $f_-(x)$ in (2.4).

via the information on two control points $P(x^-, y^-), Q(x^+, y^+)$. Note that in the classical Zadeh's summation notation (see, for instance, Chapter 2 of [27]), the above information gives $\tilde{x}_0 = 0/(x_0 - d^-) + y^-/x^- + 1/x_0 + y^+/x^+ + 0/(x_0 + d^+)$.

In the sequel, we will use the new notation $\langle x_0; d^-, d^+, \mu^-, \mu^+\rangle$ for the fuzzy number \tilde{x}_0 with the membership function (2.3)-(2.4). The advantage of the revision is that the assumption $a < b < c$ is replaced by the positivity of d^- and d^+ . We emphasize that in (2.6), x_0 represents the crisp part of the fuzzy number \tilde{x}_0 and the other four parameters represent its fuzzy part, respectively. More precisely, the fuzzy number \tilde{x}_0 can be identified by three parts: Value x_0 , Ambiguity d^-, d^+ and Fuzziness μ^-, μ^+ (see [4, 5, 21] and references therein). They capture the relevant information contained in the fuzzy number and could be useful for developing ranking procedure of fuzzy numbers. For instance, Value x_0 has the key property of permitting us to associate a real value with the fuzzy number \tilde{x}_0 . This approach is beyond the scope of this paper and will be considered elsewhere.

3 Arithmetic operators on $X_{h,p}(\mathbb{R})$

We now define the addition, multiplication, subtraction and scalar multiplication on the Gaussian-PDMF space $X_{h,p}(\mathbb{R})$ in (2.5). For convenience we shall generally use the notation \tilde{x}_i instead of $\tilde{x}_i = \langle x_i; d_i^-, d_i^+, \mu_i^-, \mu_i^+\rangle$, for any $i = 0, 1, 2, \dots$.

Definition 3.1 Let \tilde{x}_1, \tilde{x}_2 be two Gaussian-PDMFs in $X_{h,p}(\mathbb{R})$, we define

- (1) $\tilde{x}_1 \oplus \tilde{x}_2 = \langle x_1 + x_2; d_1^- d_2^-, d_1^+ d_2^+, \mu_1^- + \mu_2^-, \mu_1^+ + \mu_2^+ \rangle,$
- (2) $\lambda \tilde{x} = \langle \lambda x; (d^-)^\lambda, (d^+)^\lambda, \lambda \mu^-, \lambda \mu^+ \rangle,$
- (3) $\tilde{x}_1 \ominus \tilde{x}_2 = \langle x_1 - x_2; d_1^- (d_2^-)^{-1}, d_1^+ (d_2^+)^{-1}, \mu_1^- - \mu_2^-, \mu_1^+ - \mu_2^+ \rangle,$
- (4) $\tilde{x}_1 \otimes \tilde{x}_2 = \langle x_1 x_2; (d_1^-)^{\ln d_2^-}, (d_1^+)^{\ln d_2^+}, \mu_1^- \mu_2^-, \mu_1^+ \mu_2^+ \rangle.$

Some remarks are in order.

Remark 3.1 Note that $a < b < c$ implies $d^-, d^+ \neq 0$. Hence the crisp number $x \in \mathbb{R}$ is excluded in the Gaussian-PDMF space since it coincides to the class of vectors $\langle x; 0, 0, \cdot, \cdot \rangle$ where $d^- = d^+ = 0$. In this case, each crisp number is formally equivalent to a class

$$\{\tilde{y} = \langle x; 0, 0, \mu_y^-, \mu_y^+ \rangle, \forall \mu_y^-, \mu_y^+ \in \mathbb{R}\}.$$

Consequently, $x \in \mathbb{R}$ can be seen as the “limitation” of some quotient class of $X_{h,p}(\mathbb{R})$ satisfying

$$x \sim \lim_{d^-, d^+ \rightarrow 0} \{\tilde{x} \in X_{h,p}(\mathbb{R}) \text{ for any } \mu_y^-, \mu_y^+ \in \mathbb{R}\}.$$

and all arithmetic operations on the Gaussian-PDMF space will coincide with the corresponding operations on \mathbb{R} .

Remark 3.2 Note that in the existing literature, ranking fuzzy numbers are one of the most important research topics in fuzzy set theory. Mathematically speaking, ranking fuzzy numbers can be seen as ranking the functions as the form of (2.3) in Gaussian-PDMF space. One simple ranking method to evaluate Gaussian-PDMF numbers is that: for any $\tilde{x}, \tilde{y} \in X_{h,p}(\mathbb{R})$, $\tilde{x} \leq \tilde{y}$ if and only if $x \leq y$. As shown later, all properties of crisp numbers continues to keep under Definition 3.1.

Remark 3.3 One of the main advantage of Definition 3.1 is that, as we show later, the nice algebraic structure of Gaussian-PDMF makes the computation to be universal. For instance, it is no need to add the assumptions that the fuzzy numbers involved have to has same signs, nor the existence of the fuzzy number of H/gH difference. Furthermore, it is not necessary to use the α -cuts of the fuzzy numbers to perform calculations.

The first Theorem on the algebraic structure of Gaussian-PDMF space $X_{h,p}(\mathbb{R})$ under Definition 3.1 is:

Theorem 3.1 $X_{h,p}(\mathbb{R})$ together with the binary operation \oplus is an **abelian group**. More precisely,

(i) The binary operation \oplus is **associative**:

$$(\tilde{x}_1 \oplus \tilde{x}_2) \oplus \tilde{x}_3 = \tilde{x}_1 \oplus (\tilde{x}_2 \oplus \tilde{x}_3), \quad \text{for all } \tilde{x}_i \in X_{h,p}(\mathbb{R}), \quad i = 1, 2, 3. \quad (3.1)$$

(ii) There exists a unique **two-sided identity element** $\tilde{0} = \langle 0; 1, 1, 0, 0 \rangle$ such that

$$\tilde{0} \oplus \tilde{x} = \tilde{x} \oplus \tilde{0}, \quad \text{for all } \tilde{x} \in X_{h,p}(\mathbb{R}). \quad (3.2)$$

(iii) For every $\tilde{x} \in X_{h,p}(\mathbb{R})$ there exists a **two-sided inverse element** $-\tilde{x} \in X_{h,p}(\mathbb{R})$ such that

$$-\tilde{x} \oplus \tilde{x} = \tilde{x} \oplus -\tilde{x} = \tilde{0}. \quad (3.3)$$

Moreover, $-\tilde{x} = \langle -x; (d^-)^{-1}, (d^+)^{-1}, -\mu^-, -\mu^+ \rangle$ coincides to the scalar multiplication $\lambda \tilde{x}$ with $\lambda = -1$.

(iv) The binary operation \oplus is **commutative**:

$$\tilde{x}_1 \oplus \tilde{x}_2 = \tilde{x}_2 \oplus \tilde{x}_1, \quad \text{for all } \tilde{x}_i \in X_{h,p}(\mathbb{R}), \quad i = 1, 2. \quad (3.4)$$

Remark 3.4 Compare to the Hukuhara difference and generalized-Hukuhara difference (see, for instance, [14, 23] and references therein), the difference \ominus in Definition 3.1 shows a clear and simple algebraic structure of the fuzzy numbers. Hence, the subtraction of \tilde{x} and the addition of its negative $-\tilde{x}$ are compatible, which is not true for most of the existing definitions.

Proof of Theorem 3.1: For (i), set $\tilde{x}_i = \langle x_i; d_i^-, d_i^+, \mu_i^-, \mu_i^+ \rangle, i = 1, 2, 3$. Recall Formula (1) of Definition 3.1, it holds

$$\begin{aligned} (\tilde{x}_1 \oplus \tilde{x}_2) \oplus \tilde{x}_3 &= \langle x_1 + x_2; d_1^- d_2^-, d_1^+ d_2^+, \mu_1^- + \mu_2^-, \mu_1^+ + \mu_2^+ \rangle \oplus \langle x_3; d_3^-, d_3^+, \mu_3^-, \mu_3^+ \rangle \\ &= \langle x_1 + x_2 + x_3; d_1^- d_2^- d_3^-, d_1^+ d_2^+ d_3^+, \mu_1^- + \mu_2^- + \mu_3^-, \mu_1^+ + \mu_2^+ + \mu_3^+ \rangle \\ &= \langle x_1 + (x_2 + x_3); d_1^- (d_2^- d_3^-), d_1^+ (d_2^+ d_3^+), \mu_1^- + (\mu_2^- + \mu_3^-), \mu_1^+ + (\mu_2^+ + \mu_3^+) \rangle \\ &= \tilde{x}_1 \oplus (\tilde{x}_2 \oplus \tilde{x}_3). \end{aligned}$$

For (ii), it is trivial that $\tilde{0} = \langle 0; 1, 1, 0, 0 \rangle$ is a two-sided identity. If $\tilde{0}'$ is also a two-sided identity, then $\tilde{0} = \tilde{0} \oplus \tilde{0}' = \tilde{0}'$, which means that $\tilde{0}$ is unique.

Assertion (iii) is straightforward since for any $d^-, d^+ \neq 0$, it holds

$$-\tilde{x} \oplus \tilde{x} = \langle -x; (d^-)^{-1}, (d^+)^{-1}, -\mu^-, -\mu^+ \rangle \oplus \langle x; d^-, d^+, \mu^-, \mu^+ \rangle = \langle 0; 1, 1, 0, 0 \rangle.$$

Clearly, due to Formula (2) of Definition 3.1, $(-1)\tilde{x} = -\tilde{x}$.

Finally, the commutativity of \oplus is a direct consequence of Formula (1) of Definition 3.1 due to the fact that $d_1^- d_2^- = d_2^- d_1^-$ and $d_1^+ d_2^+ = d_2^+ d_1^+$. \square

As a direct consequence of Theorem 3.1, the linear fuzzy equation can be solved with the following formula:

Proposition 3.1 Let $\tilde{b}, \tilde{c} \in X_{h,p}(\mathbb{R})$ with $\tilde{b} = \langle b; d_b^-, d_b^+, \mu_b^-, \mu_b^+ \rangle$ and $\tilde{c} = \langle c; d_c^-, d_c^+, \mu_c^-, \mu_c^+ \rangle$. Let $a \in \mathbb{R} \setminus \{0\}$. Set $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$ be an undetermined fuzzy number such that

$$a\tilde{x} \oplus \tilde{b} = \tilde{c}. \quad (3.5)$$

Then there exists a unique fuzzy number \tilde{x} belonging to $X_{h,p}(\mathbb{R})$ satisfying (3.5) with the formula

$$\tilde{x} = \frac{1}{a}(\tilde{c} \ominus \tilde{b}) = \left\langle \frac{c-b}{a}; \left(\frac{d_c^-}{d_b^-}\right)^{\frac{1}{a}}, \left(\frac{d_c^+}{d_b^+}\right)^{\frac{1}{a}}, \frac{\mu_c^- - \mu_b^-}{a}, \frac{\mu_c^+ - \mu_b^+}{a} \right\rangle. \quad (3.6)$$

In the next theorem, we show that $X_{h,p}(\mathbb{R})$ is a commutative ring together with the operations addition and multiplication.

Theorem 3.2 $X_{h,p}(\mathbb{R})$ together with \oplus and \otimes is a **commutative ring with identity** $\tilde{1}_X = \langle 1; e, e, 1, 1 \rangle$. More precisely,

- (i) $(X_{h,p}(\mathbb{R}), \oplus)$ is an abelian group;
- (ii) $(\tilde{x}_1 \otimes \tilde{x}_2) \otimes \tilde{x}_3 = \tilde{x}_1 \otimes (\tilde{x}_2 \otimes \tilde{x}_3)$ for all $\tilde{x}_i \in X_{h,p}(\mathbb{R})$, $i = 1, 2, 3$ (**associative multiplication**);
- (iii) $\tilde{x}_1 \otimes (\tilde{x}_2 \oplus \tilde{x}_3) = \tilde{x}_1 \otimes \tilde{x}_2 \oplus \tilde{x}_1 \otimes \tilde{x}_3$ and $(\tilde{x}_1 \oplus \tilde{x}_2) \otimes \tilde{x}_3 = \tilde{x}_1 \otimes \tilde{x}_3 \oplus \tilde{x}_2 \otimes \tilde{x}_3$ for all $\tilde{x}_i \in X_{h,p}(\mathbb{R})$, $i = 1, 2, 3$ (**left and right distributive laws**);
- (iv) $\tilde{x}_1 \otimes \tilde{x}_2 = \tilde{x}_2 \otimes \tilde{x}_1$ for all $\tilde{x}_i \in X_{h,p}(\mathbb{R})$, $i = 1, 2$ (**commutative**);
- (v) $X_{h,p}(\mathbb{R})$ contains an identity element $\tilde{1}_X = \langle 1; e, e, 1, 1 \rangle$ such that

$$\tilde{1}_X \otimes \tilde{x} = \tilde{x} \otimes \tilde{1}_X \quad \text{for all } \tilde{x} \in X_{h,p}(\mathbb{R}). \quad (3.7)$$

Proof: Set $\tilde{x}_i = \langle x_i; d_i^-, d_i^+, \mu_i^-, \mu_i^+ \rangle$, $i = 1, 2, 3$. We give the proof one by one.

(i) This is Theorem 3.1.

(ii) Recall Formula (4) of Definition 3.1, it holds

$$\begin{aligned} (\tilde{x}_1 \otimes \tilde{x}_2) \otimes \tilde{x}_3 &= \langle x_1 x_2; (d_1^-)^{\ln d_2^-}, (d_1^+)^{\ln d_2^+}, \mu_1^- \mu_2^-, \mu_1^+ \mu_2^+ \rangle \otimes \langle x_3; d_3^-, d_3^+, \mu_3^-, \mu_3^+ \rangle \\ &= \left\langle x_1 x_2 x_3; (d_1^-)^{\ln d_2^- \ln d_3^-}, (d_1^+)^{\ln d_2^+ \ln d_3^+}, \mu_1^- \mu_2^- \mu_3^-, \mu_1^+ \mu_2^+ \mu_3^+ \right\rangle \\ &= \left\langle x_1 x_2 x_3; (d_1^-)^{\ln[(d_2^-)^{\ln d_3^-}]}, (d_1^+)^{\ln[(d_2^+)^{\ln d_3^+}]}, \mu_1^- \mu_2^- \mu_3^-, \mu_1^+ \mu_2^+ \mu_3^+ \right\rangle \\ &= \tilde{x}_1 \otimes (\tilde{x}_2 \otimes \tilde{x}_3). \end{aligned}$$

(iii) Recall Formula (1) and (4) of Definition 3.1, we first compute the left distributive law:

$$\begin{aligned}
& \tilde{x}_1 \otimes (\tilde{x}_2 \oplus \tilde{x}_3) \\
&= \tilde{x}_1 \otimes \langle x_2 + x_3; d_2^- + d_3^-, d_2^+ + d_3^+, \mu_2^- + \mu_3^-, \mu_2^+ + \mu_3^+ \rangle \\
&= \langle x_1(x_2 + x_3); (d_1^-)^{\ln(d_2^- d_3^-)}, (d_1^+)^{\ln(d_2^+ d_3^+)}, \mu_1^-(\mu_2^- + \mu_3^-), \mu_1^+(\mu_2^+ + \mu_3^+) \rangle \\
&= \langle x_1 x_2 + x_1 x_3; (d_1^-)^{\ln d_2^-} (d_1^-)^{\ln d_3^-}, (d_1^+)^{\ln d_2^+} (d_1^+)^{\ln d_3^+}, \mu_1^- \mu_2^- + \mu_1^- \mu_3^-, \mu_1^+ \mu_2^+ + \mu_1^+ \mu_3^+ \rangle \\
&= \tilde{x}_1 \otimes \tilde{x}_2 \oplus \tilde{x}_1 \otimes \tilde{x}_3.
\end{aligned}$$

The right distributive laws can be verified similarly.

(iv) Formula (4) of Definition 3.1 gives

$$\begin{aligned}
\tilde{x}_1 \otimes \tilde{x}_2 &= \langle x_1 x_2; (d_1^-)^{\ln d_2^-}, (d_1^+)^{\ln d_2^+}, \mu_1^- \mu_2^-, \mu_1^+ \mu_2^+ \rangle \\
&= \langle x_2 x_1; (d_2^-)^{\ln d_1^-}, (d_2^+)^{\ln d_1^+}, \mu_2^- \mu_1^-, \mu_2^+ \mu_1^+ \rangle \\
&= \tilde{x}_2 \otimes \tilde{x}_1.
\end{aligned}$$

Here we use the fact that $a^{\ln b} = b^{\ln a}$ for all $a, b > 0$.

(v) For any $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$, it holds that

$$\langle 1; e, e, 1, 1 \rangle \otimes \langle x; d^-, d^+, \mu^-, \mu^+ \rangle = \langle 1x; e^{\ln d^-}, e^{\ln d^+}, 1\mu^-, 1\mu^+ \rangle = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$$

and the second identity holds since $a^{\ln e} = a$ for any $a \in \mathbb{R}^+$.

Now we extend the binary operations on finite arithmetic operations with the following notations.

The **standard n sum** $\bigoplus_{i=1}^n \tilde{x}_i$ of $\tilde{x}_1, \dots, \tilde{x}_n \in X_{h,p}(\mathbb{R})$ is defined inductively by

$$\bigoplus_{i=1}^1 \tilde{x}_1 = \tilde{x}_1; \text{ and for } n > 1, \left(\bigoplus_{i=1}^{n-1} \tilde{x}_i \right) \oplus \tilde{x}_n$$

for any positive integer n .

Consequently, the following properties of ring hold:

Theorem 3.3 $(X_{h,p}(\mathbb{R}), \oplus, \otimes)$ has the following properties: (Properties of ring)

(i) $\tilde{0} \otimes \tilde{x} = \tilde{x} \otimes \tilde{0} = \tilde{0}$ for all $\tilde{x} \in X_{h,p}(\mathbb{R})$;

(ii) $(-\tilde{x}_1) \otimes \tilde{x}_2 = \tilde{x}_1 \otimes (-\tilde{x}_2) = -(\tilde{x}_1 \otimes \tilde{x}_2)$ for all $\tilde{x}_i \in X_{h,p}(\mathbb{R}), i = 1, 2$;

(iii) $(-\tilde{x}_1) \otimes (-\tilde{x}_2) = \tilde{x}_1 \otimes \tilde{x}_2$ for all $\tilde{x}_i \in X_{h,p}(\mathbb{R}), i = 1, 2$;

(iv) $(\lambda\tilde{x}_1) \otimes \tilde{x}_2 = \tilde{x}_1 \otimes (\lambda\tilde{x}_2) = \lambda(\tilde{x}_1 \otimes \tilde{x}_2)$ for all $\lambda \in \mathbb{R}$ and $\tilde{x}_i \in X_{h,p}(\mathbb{R}), i = 1, 2$;

(v) $\left(\bigoplus_{i=1}^n \tilde{x}_i \right) \otimes \left(\bigoplus_{j=1}^m \tilde{y}_j \right) = \bigoplus_{i=1}^n \bigoplus_{j=1}^m (\tilde{x}_i \otimes \tilde{y}_j)$ for all $\tilde{x}_i, \tilde{y}_j \in X_{h,p}(\mathbb{R}), i = 1, \dots, n, j = 1, \dots, m$.

Proof: We give the proof one by one.

(i) For any $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$, it holds that

$$\langle 0; 1, 1, 0, 0 \rangle \otimes \langle x; d^-, d^+, \mu^-, \mu^+ \rangle = \langle 0x; 1^{\ln d^-}, 1^{\ln d^+}, 0\mu^-, 0\mu^+ \rangle = \langle 0; 1, 1, 0, 0 \rangle$$

and the second identity holds since $a^{\ln 1} = 1$ for any $a \in \mathbb{R}^+$.

(ii) By the definition of \otimes , we compute

$$\begin{aligned} (-\tilde{x}_1) \otimes \tilde{x}_2 &= \langle -x_1; (d_1^-)^{-1}, (d_1^+)^{-1}, -\mu_1^-, -\mu_1^+ \rangle \otimes \langle x_2; d_2^-, d_2^+, \mu_2^-, \mu_2^+ \rangle \\ &= \langle -x_1x_2; ((d_1^-)^{-1})^{\ln d_2^-}, ((d_1^+)^{-1})^{\ln d_2^+}, -\mu_1^- \mu_2^-, -\mu_1^+ \mu_2^+ \rangle \\ &= \langle x_1(-x_2); (d_1^-)^{-\ln d_2^-}, (d_1^+)^{-\ln d_2^+}, \mu_1^-(-\mu_2^-), \mu_1^+(-\mu_2^+) \rangle \\ &= \tilde{x}_1 \otimes (-\tilde{x}_2) \end{aligned}$$

and

$$\begin{aligned} (-\tilde{x}_1) \otimes \tilde{x}_2 &= \langle -(x_1x_2); ((d_1^-)^{\ln d_2^-})^{-1}, ((d_1^+)^{\ln d_2^+})^{-1}, -(\mu_1^- \mu_2^-), -(\mu_1^+ \mu_2^+) \rangle \\ &= -(\tilde{x}_1 \otimes \tilde{x}_2). \end{aligned}$$

(iii) We compute

$$\begin{aligned} (-\tilde{x}_1) \otimes (-\tilde{x}_2) &= \langle -x_1; (d_1^-)^{-1}, (d_1^+)^{-1}, -\mu_1^-, -\mu_1^+ \rangle \otimes \langle -x_2; (d_2^-)^{-1}, (d_2^+)^{-1}, -\mu_2^-, -\mu_2^+ \rangle \\ &= \langle x_1x_2; ((d_1^-)^{-1})^{\ln(d_2^-)^{-1}}, ((d_1^+)^{-1})^{\ln(d_2^+)^{-1}}, \mu_1^- \mu_2^-, \mu_1^+ \mu_2^+ \rangle \\ &= \langle x_1x_2; (d_1^-)^{\ln d_2^-}, (d_1^+)^{\ln d_2^+}, \mu_1^- \mu_2^-, \mu_1^+ \mu_2^+ \rangle \\ &= \tilde{x}_1 \otimes \tilde{x}_2. \end{aligned}$$

(iv) By the definition of scalar multiplication, we compute

$$\begin{aligned}
(\lambda \tilde{x}_1) \otimes \tilde{x}_2 &= \langle -x_1; (d_1^-)^\lambda, (d_1^+)^\lambda, \lambda \mu_1^-, \lambda \mu_1^+ \rangle \otimes \langle x_2; d_2^-, d_2^+, \mu_2^-, \mu_2^+ \rangle \\
&= \langle \lambda x_1 x_2; ((d_1^-)^\lambda)^{\ln d_2^-}, ((d_1^+)^\lambda)^{\ln d_2^+}, \lambda \mu_1^- \mu_2^-, \lambda \mu_1^+ \mu_2^+ \rangle \\
&= \langle x_1(\lambda x_2); (d_1^-)^{\lambda \ln d_2^-}, (d_1^+)^{\lambda \ln d_2^+}, \mu_1^-(\lambda \mu_2^-), \mu_1^+(\lambda \mu_2^+) \rangle \\
&= \tilde{x}_1 \otimes (\lambda \tilde{x}_2)
\end{aligned}$$

and

$$\begin{aligned}
(\lambda \tilde{x}_1) \otimes \tilde{x}_2 &= \langle \lambda(x_1 x_2); \left((d_1^-)^{\ln d_2^-} \right)^\lambda, \left((d_1^+)^{\ln d_2^+} \right)^\lambda, \lambda(\mu_1^- \mu_2^-), \lambda(\mu_1^+ \mu_2^+) \rangle \\
&= \lambda(\tilde{x}_1 \otimes \tilde{x}_2).
\end{aligned}$$

(v) We prove it by induction. The assertion is trivial for $m = n = 1$. Assume that

$$(\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_n) \otimes (\tilde{y}_1 \oplus \cdots \oplus \tilde{y}_m) = \bigoplus_{i=1}^n \bigoplus_{j=1}^m (\tilde{x}_i \otimes \tilde{y}_j)$$

holds for all $n \leq N$ and $m \leq M$. Then by the distributive law (Formula (iii) of Theorem 3.2), it holds

$$\begin{aligned}
&(\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_N \oplus \tilde{x}_{N+1}) \otimes (\tilde{y}_1 \oplus \cdots \oplus \tilde{y}_M) \\
&= (\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_N) \otimes (\tilde{y}_1 \oplus \cdots \oplus \tilde{y}_M) \oplus \tilde{x}_{N+1} \otimes (\tilde{y}_1 \oplus \cdots \oplus \tilde{y}_M) \\
&= \bigoplus_{i=1}^N \bigoplus_{j=1}^M (\tilde{x}_i \otimes \tilde{y}_j) \oplus (\tilde{x}_{N+1} \otimes \tilde{y}_1) \oplus \cdots \oplus (\tilde{x}_{N+1} \otimes \tilde{y}_M) \\
&= \bigoplus_{i=1}^{N+1} \bigoplus_{j=1}^M (\tilde{x}_i \otimes \tilde{y}_j).
\end{aligned}$$

Similarly,

$$(\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_N) \otimes (\tilde{y}_1 \oplus \cdots \oplus \tilde{y}_M \oplus \tilde{y}_{M+1}) = \bigoplus_{i=1}^N \bigoplus_{j=1}^{M+1} (\tilde{x}_i \otimes \tilde{y}_j).$$

Hence the assertion holds true for any $m, n \in \mathbb{N}$. \square

We now consider those elements in $X_{h,p}(\mathbb{R})$ which have an inverse under multiplication. They form a group, and this “group of units” is very important in algebraic structure theory. We start with the following definitions:

Definition 3.2 An element \tilde{x} in $X_{h,p}(\mathbb{R})$ with is said to be **left** (resp. **right**) **invertible** if there exists $\tilde{y} \in X_{h,p}(\mathbb{R})$ (resp. $\tilde{z} \in X_{h,p}(\mathbb{R})$) such that $\tilde{y} \otimes \tilde{x} = \tilde{1}_X$ (reps. $\tilde{x} \otimes \tilde{z} = \tilde{1}_X$). The element \tilde{y} (resp. \tilde{z}) is called a **left** (resp. **right**) **inverse** of \tilde{x} . An element $\tilde{x} \in X_{h,p}(\mathbb{R})$ that is both left and right invertible is denoted by \tilde{x}^{-1} and said to be **invertible** or to be a **unit**.

Theorem 3.4 Let

$$U(X) = \{\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle \in X_{h,p}(\mathbb{R}) \mid x \neq 0, d^- \neq 1, d^+ \neq 1, \mu^- \neq 0, \mu^+ \neq 0\}. \quad (3.8)$$

Then it is the set of all units in $X_{h,p}(\mathbb{R})$. Moreover, for any $\tilde{x} \in U(X)$,

$$\tilde{x} \otimes \tilde{y} = \tilde{0} \text{ or } \tilde{y} \otimes \tilde{x} = \tilde{0} \implies \tilde{y} = \tilde{0}. \quad (3.9)$$

Remark 3.5 Note that for $\tilde{x}_1, \tilde{x}_2 \in X_{h,p}(\mathbb{R})$, the following assertion is false:

$$\tilde{x}_1 \otimes \tilde{x}_2 = \tilde{0} \implies \tilde{x}_1 = \tilde{0} \text{ or } \tilde{x}_2 = \tilde{0}.$$

In fact, any $\tilde{x}_2 = \langle 0; d_2^-, d_2^+, \mu_2^-, \mu_2^+ \rangle$, with the leading term $x_2 = 0$, acts as both a left and right zero divisor of the nonzero element $\tilde{x}_1 = \langle x_1; 1, 1, 0, 0 \rangle$. According to the definition of the zero element $\tilde{0}$ in $X_{h,p}(\mathbb{R})$, the following two expressions are equivalent:

$$\tilde{x}_1 \otimes \tilde{x}_2 = \tilde{0} \iff x_1 x_2 = 0, \ln d_1^- \ln d_2^- = 0, \ln d_1^+ \ln d_2^+ = 0, \mu_1^- \mu_2^- = 0, \mu_1^+ \mu_2^+ = 0.$$

Consequently, all elements in $U(X)$ are invertible.

Remark 3.6 Note that the right and left cancellation laws hold in $U(X)$ by Assertion (3.9), that is, for all $\tilde{x}_i \in U(X), i = 1, 2, 3$,

$$\tilde{x}_1 \otimes \tilde{x}_2 = \tilde{x}_1 \otimes \tilde{x}_3 \text{ or } \tilde{x}_2 \otimes \tilde{x}_1 = \tilde{x}_3 \otimes \tilde{x}_1 \implies \tilde{x}_2 = \tilde{x}_3.$$

Proof of Theorem 3.4: For any $\tilde{x} \in U(X)$, taking into account that $\tilde{1}_X = \langle 1; e, e, 1, 1 \rangle$, the inverse of \tilde{x} is uniquely given by

$$\tilde{x}^{-1} = \left\langle \frac{1}{x}; \exp\left(\frac{1}{\ln d^-}\right), \exp\left(\frac{1}{\ln d^+}\right), \frac{1}{\mu^-}, \frac{1}{\mu^+} \right\rangle$$

Clearly, (3.9) is trivial. □

As a direct consequence of Theorem 3.4, it is possible to update the result for the first order fuzzy linear equation (3.5) in Proposition 3.1. We have

Proposition 3.2 Let $\tilde{a} = \langle a; d_a^-, d_a^+, \mu_a^-, \mu_a^+ \rangle$ be in $U(X)$, $\tilde{b} = \langle b; d_b^-, d_b^+, \mu_b^-, \mu_b^+ \rangle$ and $\tilde{c} = \langle c; d_c^-, d_c^+, \mu_c^-, \mu_c^+ \rangle$ be in $X_{h,p}(\mathbb{R})$. Set $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$ be an undetermined fuzzy number such that

$$\tilde{a}\tilde{x} \oplus \tilde{b} = \tilde{c}. \quad (3.10)$$

Then there exists a unique fuzzy number \tilde{x} belonging to $X_{h,p}(\mathbb{R})$ satisfying (3.10). More precisely,

$$\tilde{x} = \left\langle \frac{c-b}{a}; \exp\left(\frac{\ln d_c^- - \ln d_b^-}{\ln d_a^-}\right), \exp\left(\frac{\ln d_c^+ - \ln d_b^+}{\ln d_a^+}\right), \frac{\mu_c^- - \mu_b^-}{\mu_a^-}, \frac{\mu_c^+ - \mu_b^+}{\mu_a^+} \right\rangle. \quad (3.11)$$

The next binomial theorem is frequently useful in computations. We define \tilde{x}^0 be the identity element $\tilde{1}_X$. The element \tilde{x}^n is defined to be the standard n product $\tilde{x} \otimes \tilde{x} \otimes \cdots \otimes \tilde{x}$. Recall that if k and n are integers with $0 \leq k \leq n$, then the **binomial coefficient** $\binom{n}{k}$ is the number $n!/(n-k)!k!$, where $0! = 1$ and $n! = n(n-1)(n-2)\cdots 2 \cdot 1$ for $n \geq 1$. It is obvious that $\binom{n}{k}$ is actually an integer.

Theorem 3.5 (Binomial Theorem). *Let n be a positive integer and $\tilde{x}, \tilde{y}, \tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_n \in X_{h,p}(\mathbb{R})$.*

$$(i) \quad (\tilde{x} \oplus \tilde{y})^n = \bigoplus_{k=0}^n \binom{n}{k} \tilde{x}^k \otimes \tilde{y}^{n-k};$$

(ii) *Moreover,*

$$(\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_s)^n = \bigoplus \frac{n!}{(i_1!) \cdots (i_s!)} \tilde{x}_1^{i_1} \otimes \tilde{x}_2^{i_2} \cdots \otimes \tilde{x}_s^{i_s},$$

where the sum is over all s -tuples (i_1, i_2, \dots, i_s) such that $i_1 + i_2 + \cdots + i_s = n$.

Proof: Assertion (i) can be proved using induction on n and the fact that \tilde{x} and \tilde{y} satisfy the distributive law and the commutativity. In fact, we compute

$$\begin{aligned} (\tilde{x} \oplus \tilde{y})^{n+1} &= (\tilde{x} \oplus \tilde{y})^n \otimes (\tilde{x} \oplus \tilde{y}) \\ &= \left(\bigoplus_{k=0}^n \binom{n}{k} \tilde{x}^k \otimes \tilde{y}^{n-k} \otimes \tilde{x} \right) \oplus \left(\bigoplus_{k=0}^n \binom{n}{k} \tilde{x}^k \otimes \tilde{y}^{n-k} \otimes \tilde{y} \right) \\ &= \tilde{x}^{n+1} \oplus \left(\bigoplus_{k=0}^{n-1} \binom{n}{k} \tilde{x}^{k+1} \otimes \tilde{y}^{n-k} \right) \oplus \left(\bigoplus_{k=1}^n \binom{n}{k} \tilde{x}^k \otimes \tilde{y}^{n-k+1} \oplus \tilde{y}^{n+1} \right) \oplus \tilde{y}^{n+1} \\ &= \tilde{x}^{n+1} \oplus \left(\bigoplus_{k=0}^{n-1} \left(\binom{n}{k} + \binom{n}{k+1} \right) \tilde{x}^{k+1} \otimes \tilde{y}^{n-k} \right) \oplus \tilde{y}^{n+1} \\ &= \tilde{x}^{n+1} \oplus \left(\bigoplus_{k=0}^{n-1} \binom{n+1}{k+1} \tilde{x}^{k+1} \otimes \tilde{y}^{n-k} \right) \oplus \tilde{y}^{n+1} \\ &= \tilde{x}^{n+1} \oplus \left(\bigoplus_{k=1}^n \binom{n+1}{k} \tilde{x}^k \otimes \tilde{y}^{n+1-k} \right) \oplus \tilde{y}^{n+1} \\ &= \bigoplus_{k=0}^{n+1} \binom{n+1}{k} \tilde{x}^k \otimes \tilde{y}^{n+1-k}. \end{aligned}$$

Note that we use the formula that for $k < n$, it holds

$$\begin{aligned}
\binom{n}{k} + \binom{n}{k+1} &= \frac{n!}{(n-k)!k!} + \frac{n!}{(n-k-1)!(k+1)!} \\
&= \frac{n!(k+1)}{(n-k)!(k+1)!} + \frac{n!(n-k)}{(n-k)!(k+1)!} \\
&= \frac{n!(n+1)}{(n-k)!(k+1)!} = \binom{n+1}{k+1}.
\end{aligned}$$

Assertion (ii) can be proved using induction on s . The case $s = 2$ is just part (i) since

$$(\tilde{x}_1 \oplus \tilde{x}_2)^n = \bigoplus_{k=0}^n \binom{n}{k} \tilde{x}_1^k \otimes \tilde{x}_2^{n-k} = \bigoplus_{k+j=n} \frac{n!}{k!j!} \tilde{x}_1^k \otimes \tilde{x}_2^j.$$

Assume that (ii) is true for s . Note that

$$\begin{aligned}
(\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_s \oplus \tilde{x}_{s+1})^n &= ((\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_s) \oplus \tilde{x}_{s+1})^n \\
&= \bigoplus_{k=0}^n \binom{n}{k} (\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_s)^k \otimes \tilde{x}_{s+1}^{n-k} \\
&= \bigoplus_{k+j=n} \frac{n!}{k!j!} (\tilde{x}_1 \oplus \cdots \oplus \tilde{x}_s)^k \otimes \tilde{x}_{s+1}^j \\
&= \bigoplus_{k+j=n} \frac{n!}{k!j!} \left(\bigoplus_{i_1+i_2+\cdots+i_s=k} \frac{k!}{(i_1!) \cdots (i_s!)} \tilde{x}_1^{i_1} \otimes \tilde{x}_2^{i_2} \cdots \otimes \tilde{x}_s^{i_s} \right) \otimes \tilde{x}_{s+1}^j \\
&= \bigoplus_{i_1+i_2+\cdots+i_s+j=n} \frac{n!}{(i_1!) \cdots (i_s!)(j!)} \tilde{x}_1^{i_1} \otimes \tilde{x}_2^{i_2} \cdots \otimes \tilde{x}_s^{i_s} \otimes \tilde{x}_{s+1}^j
\end{aligned}$$

by part (i). The proof is complete. \square

As a simple consequence of Theorem 3.5 with $n = 2$, we solve the following fuzzy quadratic equation.

Proposition 3.3 *Let $\tilde{b}, \tilde{c} \in X_{h,p}(\mathbb{R})$ with $\tilde{b} = \langle b; d_b^-, d_b^+, \mu_b^-, \mu_b^+ \rangle$ and $\tilde{c} = \langle c; d_c^-, d_c^+, \mu_c^-, \mu_c^+ \rangle$. Set $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$ be an undetermined fuzzy number such that*

$$\tilde{x}^2 \ominus (\tilde{b} \oplus \tilde{c}) \otimes \tilde{x} \oplus (\tilde{b} \otimes \tilde{c}) = \tilde{0}. \quad (3.12)$$

Then $\tilde{x} = \tilde{b}$ and $\tilde{x} = \tilde{c}$ are solutions of the equation (3.12).

Remark 3.7 *Note that the solvability of fuzzy quadratic equation is much more complicated than the standard quadratic equation due to the lack of cancellation law in $X_{h,p}(\mathbb{R})$ (see Remark (3.5)). Nevertheless, under some suitable conditions on the coefficients $\tilde{a}, \tilde{b}, \tilde{c}$, one can give an quadratic formula for the fuzzy quadratic polynomial*

$$\tilde{a}\tilde{x}^2 \oplus \tilde{b}\tilde{x} \oplus \tilde{c} = \tilde{0}. \quad (3.13)$$

One possible assumption is that $\tilde{a} \in U(X)$ and Equation (3.13) can be expressed as a product of

$$(\tilde{p}\tilde{x} \oplus \tilde{q}) \otimes (\tilde{r}\tilde{x} \oplus \tilde{s}) = \tilde{0}.$$

Moreover, one can expect that the roots \tilde{x}_1, \tilde{x}_2 of the fuzzy quadratic polynomial $P(\tilde{x}) = \tilde{a}\tilde{x}^2 \oplus \tilde{b}\tilde{x} \oplus \tilde{c}$ satisfy

$$\tilde{x}_1 \oplus \tilde{x}_2 = \ominus \tilde{a}^{-1} \tilde{b}, \quad \tilde{x}_1 \otimes \tilde{x}_2 = \tilde{a}^{-1} \tilde{c},$$

which is the **fuzzy Vieta's formulas** for $\tilde{a} \in U(X)$. Compare to the “crisp” polynomial case, the main difference is that the fuzzy numbers are quinary arrays and a systematic analysis of the algebraic structure is needed.

4 Vector space

We analyze the Gaussian-PDMF space with two operations: **addition and scalar multiplication**. We claim that,

Theorem 4.1 $X_{h,p}(\mathbb{R})$ is a vector space satisfying the following ten properties (axioms):

- (1) The sum of \tilde{x}_1, \tilde{x}_2 , denoted by $\tilde{x}_1 \oplus \tilde{x}_2$, is in $X_{h,p}(\mathbb{R})$.
- (2) $\tilde{x}_1 \oplus \tilde{x}_2 = \tilde{x}_2 \oplus \tilde{x}_1$.
- (3) $(\tilde{x}_1 \oplus \tilde{x}_2) \oplus \tilde{x}_3 = \tilde{x}_1 \oplus (\tilde{x}_2 \oplus \tilde{x}_3)$.
- (4) There is a **zero** vector $\tilde{0}$ in $X_{h,p}(\mathbb{R})$ such that $\tilde{x} \oplus \tilde{0} = \tilde{x}$ for any $\tilde{x} \in X_{h,p}(\mathbb{R})$.
- (5) For each $\tilde{x} \in X_{h,p}(\mathbb{R})$, there is a vector $-\tilde{x}$ in $X_{h,p}(\mathbb{R})$ such that $\tilde{x} \oplus (-\tilde{x}) = \tilde{0}$.
- (6) The scalar multiplication of \tilde{x} by λ , denoted by $\lambda\tilde{x}$, is in $X_{h,p}(\mathbb{R})$.
- (7) $\lambda(\tilde{x}_1 \oplus \tilde{x}_2) = \lambda\tilde{x}_1 \oplus \lambda\tilde{x}_2$.
- (8) $(\lambda_1 + \lambda_2)\tilde{x} = \lambda_1\tilde{x} \oplus \lambda_2\tilde{x}$.
- (9) $\lambda_1(\lambda_2\tilde{x}) = (\lambda_1\lambda_2)\tilde{x}$.
- (10) $1\tilde{x} = \tilde{x}$.

Remark 4.1 Technically, $X_{h,p}(\mathbb{R})$ is a real vector space. All of the theory in this section should hold for a complex vector space if the scalars are complex numbers and Rule (2) of Definition 3.1 is adapted to complex numbers. The detail of the proof is beyond the scope of this paper and need to be done in the future.

Proof of Theorem 4.1: Assertion (1) – (3), (6) are trivial. (4) and (5) is the assertion (ii) and (iii) of Theorem 3.1.

For (7) – (10), by Formula (1) and (2) of (3.1), we compute

$$\begin{aligned}
\lambda(\tilde{x}_1 \oplus \tilde{x}_2) &= \lambda(\langle x_1; d_1^-, d_1^+, \mu_1^-, \mu_1^+ \rangle \oplus \langle x_2; d_2^-, d_2^+, \mu_2^-, \mu_2^+ \rangle) \\
&= \lambda(\langle x_1 + x_2; d_1^- d_2^-, d_1^+ d_2^+, \mu_1^- \mu_2^-, \mu_1^+ \mu_2^+ \rangle) \\
&= \langle \lambda(x_1 + x_2); (d_1^- d_2^-)^\lambda, (d_1^+ d_2^+)^\lambda, \lambda(\mu_1^- + \mu_2^-), \lambda(\mu_1^+ + \mu_2^+) \rangle \\
&= \langle \lambda x_1 + \lambda x_2; (d_1^-)^\lambda (d_2^-)^\lambda, (d_1^+)^\lambda (d_2^+)^\lambda, \lambda \mu_1^- + \lambda \mu_2^-, \lambda \mu_1^+ + \lambda \mu_2^+ \rangle \\
&= \lambda \tilde{x}_1 \oplus \lambda \tilde{x}_2,
\end{aligned}$$

and

$$\begin{aligned}
(\lambda_1 + \lambda_2)\tilde{x} &= \langle (\lambda_1 + \lambda_2)x; (d^-)^{\lambda_1 + \lambda_2}, (d^+)^{\lambda_1 + \lambda_2}, (\lambda_1 + \lambda_2)\mu^-, (\lambda_1 + \lambda_2)\mu^+ \rangle \\
&= \langle \lambda_1 x; (d^-)^{\lambda_1}, (d^+)^{\lambda_1}, \lambda_1 \mu^-, \lambda_1 \mu^+ \rangle \oplus \langle \lambda_2 x; (d^-)^{\lambda_2}, (d^+)^{\lambda_2}, \lambda_2 \mu^-, \lambda_2 \mu^+ \rangle \\
&= \lambda_1 \tilde{x} \oplus \lambda_2 \tilde{x},
\end{aligned}$$

and

$$\begin{aligned}
\lambda_1(\lambda_2 \tilde{x}) &= \lambda_1 \langle \lambda_2 x; (d^-)^{\lambda_2}, (d^+)^{\lambda_2}, \lambda_2 \mu^-, \lambda_2 \mu^+ \rangle \\
&= \langle \lambda_1 \lambda_2 x; ((d^-)^{\lambda_2})^{\lambda_1}, ((d^+)^{\lambda_2})^{\lambda_1}, \lambda_1 \lambda_2 \mu^-, \lambda_1 \lambda_2 \mu^+ \rangle \\
&= (\lambda_1 \lambda_2) \tilde{x},
\end{aligned}$$

and finally

$$1\tilde{x} = \langle 1x; (d_1^-)^1, (d_1^+)^1, 1\mu_1^-, 1\mu_1^+ \rangle = \tilde{x}.$$

□

In the sequel, we present a basis of the linear space $X_{h,p}(\mathbb{R})$.

Definition 4.1 *An indexed set of vectors $\mathcal{X} = \{\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_p\}$ in $X_{h,p}(\mathbb{R})$ is said to be linearly independent if the equation*

$$\lambda_1 \tilde{x}_1 \oplus \lambda_2 \tilde{x}_2 \oplus \dots \oplus \lambda_p \tilde{x}_p = \tilde{0}$$

has only the trivial solution, $\lambda_1 = 0, \dots, \lambda_p = 0$.

Under the above definition, we have

Theorem 4.2 A set of vectors $\mathcal{X} = \{\tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{x}_4, \tilde{x}_5\}$ in $X_{h,p}(\mathbb{R})$ as the form

$$\tilde{x}_1 = \langle 1; 1, 1, 0, 0 \rangle, \tilde{x}_2 = \langle 0; e, 1, 0, 0 \rangle, \tilde{x}_3 = \langle 0; 1, e, 0, 0 \rangle, \tilde{x}_4 = \langle 0; 1, 1, 1, 0 \rangle, \tilde{x}_5 = \langle 0; 1, 1, 0, 1 \rangle$$

is a **basis** for $X_{h,p}(\mathbb{R})$. More precisely,

1. \mathcal{X} is a linear independent set, and
2. $X_{h,p}(\mathbb{R})$ can be spanned by \mathcal{X} , i.e. for any $\tilde{x} \in X_{h,p}(\mathbb{R})$, there exists $\lambda_i \in \mathbb{R}, i = 1, 2, 3, 4, 5$ such that

$$\tilde{x} = \sum_{i=1}^5 \lambda_i \tilde{x}_i = \lambda_1 \tilde{x}_1 \oplus \lambda_2 \tilde{x}_2 \oplus \lambda_3 \tilde{x}_3 \oplus \lambda_4 \tilde{x}_4 \oplus \lambda_5 \tilde{x}_5.$$

Moreover, the set of scalars $\lambda_1, \dots, \lambda_5$ is unique.

We call that the **coordinates of \tilde{x} relative to the basis \mathcal{X}** (or the **\mathcal{X} -coordinates of \tilde{x}**) are the weight $\lambda_1, \dots, \lambda_5$ such that

$$\tilde{x} = \lambda_1 \tilde{x}_1 \oplus \lambda_2 \tilde{x}_2 \oplus \lambda_3 \tilde{x}_3 \oplus \lambda_4 \tilde{x}_4 \oplus \lambda_5 \tilde{x}_5.$$

We call the vector in \mathbb{R}^5

$$\left[\tilde{x} \right]_{\mathcal{X}} = \left[\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5 \right]$$

the **\mathcal{X} -coordinate vector of \tilde{x}** . The mapping $x \mapsto \left[\tilde{x} \right]_{\mathcal{X}}$ is the **coordinate mapping determined by \mathcal{X}** . In fact, it connects the Gaussian-PDMF space $X_{h,p}(\mathbb{R})$ to the standard Euclidean space \mathbb{R}^5 . We have

Theorem 4.3 The \mathcal{X} -coordinate vector of $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$ is given by

$$\left[\tilde{x} \right]_{\mathcal{X}} = \left[x, \ln d^-, \ln d^+, \mu^-, \mu^+ \right]$$

for any $\tilde{x} \in X_{h,p}(\mathbb{R})$. Moreover, the coordinate mapping $x \mapsto \left[\tilde{x} \right]_{\mathcal{X}}$ is a one-to-one linear map (isomorphism) from $X_{h,p}(\mathbb{R})$ onto \mathbb{R}^5 , i.e., for any $\lambda_1, \lambda_2 \in \mathbb{R}$,

$$\left[\lambda_1 \tilde{x}_1 \oplus \lambda_2 \tilde{x}_2 \right]_{\mathcal{X}} = \lambda_1 \left[\tilde{x}_1 \right]_{\mathcal{X}} + \lambda_2 \left[\tilde{x}_2 \right]_{\mathcal{X}}.$$

Proof: It is straightforward. □

5 Numerical simulation

In this section, we illustrate three examples to show the advantage of the Gaussian-PDMF space with the algebraic structures. The first and the second example are linear first order

fuzzy equation, with real numbers and fuzzy numbers coefficients, respectively. The third one is a quadratic monic fuzzy equation.

Example 1. Let $\tilde{b}, \tilde{c} \in X_{h,p}(\mathbb{R})$ with $\tilde{1} = \langle 1; 1, 1, 0, 0 \rangle$ and $\tilde{3} = \langle 3; 4, 1, 1, 2 \rangle$. Let $a = 2$. Set $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$ be an undetermined fuzzy number in $X_{h,p}(\mathbb{R})$ satisfying (3.5), i.e.,

$$2\tilde{x} \oplus \tilde{1} = \tilde{3}. \quad (5.1)$$

Solution A. Using Proposition 3.1, \tilde{x} is given by

$$\tilde{x} = \frac{1}{2}(\tilde{3} \ominus \tilde{1}) = \langle 1; 2, 1, \frac{1}{2}, 1 \rangle.$$

The corresponding fuzzy numbers are in Figure 2 and Figure 3.




Figure 2: The graph of $\tilde{3}$ and $\tilde{1}$




Figure 3: The graph of $\frac{1}{2}(\tilde{3} \ominus \tilde{1})$

Solution B. Using Theorem 4.3, we compute the \mathcal{X} -coordinate vector of \tilde{x} as

$$[\tilde{1}]_{\mathcal{X}} = [1, 0, 0, 0, 0]_{\mathcal{X}}, \quad [\tilde{3}]_{\mathcal{X}} = [3, \ln 4, 0, 1, 2]_{\mathcal{X}}$$

(5.1) holds if and only if

$$2[\tilde{x}]_{\mathcal{X}} + [\tilde{1}]_{\mathcal{X}} = [\tilde{3}]_{\mathcal{X}}. \quad (5.2)$$

(5.2) is a standard arithmetic equation and can be solved as

$$[\tilde{x}]_{\mathcal{X}} = [1, \ln 2, 0, \frac{1}{2}, 1]_{\mathcal{X}},$$

It means that

$$\tilde{x} = \langle 1; 2, 1, \frac{1}{2}, 1 \rangle.$$

We show an example of Proposition 3.2, which solves $\tilde{a}\tilde{x} \oplus \tilde{b} = \tilde{c}$.

Example 2. Let $\tilde{b}, \tilde{c} \in X_{h,p}(\mathbb{R})$ with $\tilde{1} = \langle 1; 1, 1, 0, 0 \rangle$ and $\tilde{3} = \langle 3; 4, 1, 1, 2 \rangle$. Let $\tilde{a} = \langle 2; e, e, 1, 1 \rangle$. Set $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$ be an undetermined fuzzy number in $X_{h,p}(\mathbb{R})$ satisfying (3.10), i.e.,

$$\tilde{2}\tilde{x} \oplus \tilde{1} = \tilde{3}. \quad (5.3)$$

Solution. Note that Theorem 4.3 is no longer applicable and we apply Proposition 3.2. We first compute

$$\tilde{2}^{-1} = \langle \frac{1}{2}; 1, 1, 1, 1 \rangle, \quad \tilde{3} \ominus \tilde{1} = \langle 2; 4, 1, 1, 2 \rangle.$$

Hence

$$\tilde{x} = \tilde{2}^{-1} \otimes (\tilde{3} \ominus \tilde{1}) = \langle 1; 1, 1, 1, 2 \rangle.$$

The corresponding fuzzy numbers are in Figure 4 and Figure 5.




Figure 4: The graph of $\tilde{2}$




Figure 5: The graph of $\tilde{2}^{-1} \otimes (\tilde{3} \ominus \tilde{1})$

Example 3. Let $\tilde{2} = \langle 2; e, e, 1, 2 \rangle$ and $\tilde{3} = \langle 3; 1, 1, 0, 1 \rangle$. Set $\tilde{x} = \langle x; d^-, d^+, \mu^-, \mu^+ \rangle$ be an undetermined fuzzy number in $X_{h,p}(\mathbb{R})$ satisfying

$$\tilde{x}^2 \oplus \tilde{2}\tilde{x} \ominus \tilde{3} = \tilde{0}. \quad (5.4)$$

Solution. We solve the 5-tuple \tilde{x} one by one.

- For $X_{h,p}(\mathbb{R})$, we have $x^2 + 2x - 3 = 0$. Hence, $x_1 = 1$ or $x_2 = -3$;
- For d^- , we have

$$(d^-)^{\ln d^-} \cdot e^{\ln d^-} \cdot 1^{-1} = 1 \Leftrightarrow (\ln d^-)^2 + \ln d^- + 0 = 0 \Leftrightarrow \ln d^- (\ln d^- + 1) = 0.$$

Hence, $d_1^- = 1$ or $d_2^- = e^{-1}$;

- For d^+ , we have the same equation as d^- . Hence, $d_1^+ = 1$ or $d_2^+ = e^{-1}$;
- For μ^- , we have $(\mu^-)^2 + \mu^- = 0$. Hence, $\mu_1^- = -1$ or $\mu_2^- = 0$;
- For μ^+ , we have $(\mu^+)^2 + 2\mu^+ + 1 = 0$. Hence, $\mu_{1,2}^+ = -1$.

Interestingly, there are 2^4 fuzzy numbers in $X_{h,p}(\mathbb{R})$ satisfying Equation (5.4):

$$\tilde{x} = \langle x_i; d_j^-, d_k^+, \mu_l^-, -1 \rangle \text{ for any } i, j, k, l = 1, 2.$$

6 Final remarks

In summary, our research introduces a new class of membership functions, merging subjective perception and objective information through the Gaussian Probability Density Membership Function (Gaussian-PDMF) space. This space, denoted as $X_{h,p}(\mathbb{R})$, uniquely identifies fuzzy numbers, symbolized by \tilde{x} , with a vector notation $\langle x; d^-, d^+, \mu^-, \mu^+ \rangle$. We define five algebraic operations, revealing a robust structure within $X_{h,p}(\mathbb{R})$. Especially, addition \oplus forms an abelian group, and multiplication \otimes establishes a commutative ring with identity. This structured algebraic framework proves instrumental in solving fuzzy equations, applying the Binomial Theorem, and exploring fuzzy Vieta's formula.

Our work could extend theoretical insights into fuzzy numbers, providing a fresh framework for understanding and applying fuzzy mathematics. The practical utility of this algebraic structure is demonstrated through several numerical examples. We hope this research should contribute to the broader field of fuzzy mathematics and related research areas.

References

- [1] K. T. ATANASSOV, *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems, 20 (1986), pp. 87–96.
- [2] K. T. ATANASSOV, *Intuitionistic fuzzy sets*, vol. 35 of Studies in Fuzziness and Soft Computing, Physica-Verlag, Heidelberg, 1999. Theory and applications.
- [3] V. S. COSTA, B. BEDREGAL, AND R. H. SANTIAGO, *On closure properties of \mathcal{L} -valued linear languages*, Fuzzy Sets and Systems, 420 (2021), pp. 54–71.
- [4] M. DELGADO, M. A. VILA, AND W. VOXMAN, *A fuzziness measure for fuzzy numbers: Applications*, Fuzzy sets and systems, 94 (1998), pp. 205–216.
- [5] ———, *On a canonical representation of fuzzy numbers*, Fuzzy sets and systems, 93 (1998), pp. 125–135.
- [6] J. DOMBI, *Membership function as an evaluation*, Fuzzy sets and systems, 35 (1990), pp. 1–21.
- [7] D. DUBOIS AND H. PRADE, *Fuzzy sets and systems: theory and applications*, Academic press, 1980.
- [8] D. DUBOIS AND H. PRADE, *Additions of interactive fuzzy numbers*, IEEE Transactions on Automatic Control, 26 (1981), pp. 926–936.
- [9] D. DUBOIS AND H. PRADE, *Towards fuzzy differential calculus part 3: Differentiation*, Fuzzy sets and systems, 8 (1982), pp. 225–233.

- [10] E. ESMI, L. C. DE BARROS, AND V. F. WASQUES, *Some notes on the addition of interactive fuzzy numbers*, in *Fuzzy Techniques: Theory and Applications*, R. B. Kearfott, I. Batyrshin, M. Reformat, M. Ceberio, and V. Kreinovich, eds., Cham, 2019, Springer International Publishing, pp. 246–257.
- [11] E. ESMI, V. F. WASQUES, AND L. CARVALHO DE BARROS, *Addition and subtraction of interactive fuzzy numbers via family of joint possibility distributions*, *Fuzzy Sets and Systems*, 424 (2021), pp. 105–131. Uncertainty.
- [12] R. E. GIACHETTI AND R. E. YOUNG, *A parametric representation of fuzzy numbers and their arithmetic operators*, *Fuzzy sets and systems*, 91 (1997), pp. 185–202.
- [13] M. L. GUERRA AND L. STEFANINI, *Approximate fuzzy arithmetic operations using monotonic interpolations*, *Fuzzy Sets and Systems*, 150 (2005), pp. 5–33.
- [14] M. HUKUHARA, *Integration des applications mesurables dont la valeur est un compact convexe*, *Funkcialaj Ekvacioj*, 10 (1967), pp. 205–223.
- [15] E. P. KLEMENT, R. MESIAR, AND E. PAP, *Triangular norms. position paper i: basic analytical and algebraic properties*, *Fuzzy Sets and Systems*, 143 (2004), pp. 5–26. *Advances in Fuzzy Logic*.
- [16] E. KRUSIŃSKA AND J. LIEBHART, *A note on the usefulness of linguistic variables for differentiating between some respiratory diseases*, *Fuzzy sets and systems*, 18 (1986), pp. 131–142.
- [17] Y. LI AND W. PEDRYCZ, *Fuzzy finite automata and fuzzy regular expressions with membership values in lattice-ordered monoids*, *Fuzzy sets and systems*, 156 (2005), pp. 68–92.
- [18] D. LIANG, D. LIU, W. PEDRYCZ, AND P. HU, *Triangular fuzzy decision-theoretic rough sets*, *International Journal of Approximate Reasoning*, 54 (2013), pp. 1087–1106.
- [19] S. MASHCHENKO, *Sums of fuzzy sets of summands*, *Fuzzy Sets and Systems*, 417 (2021), pp. 140–151.
- [20] M. L. PURI AND D. A. RALESCU, *Differentials of fuzzy functions*, *Journal of Mathematical Analysis and Applications*, 91 (1983), pp. 552–558.
- [21] N. A. RAHMAN, L. ABDULLAH, A. T. A. GHANI, AND N. AHMAD, *Solutions of interval type-2 fuzzy polynomials using a new ranking method*, in *AIP Conference Proceedings*, vol. 1682, AIP Publishing, 2015.
- [22] Y. SHEN, *First-order linear fuzzy differential equations on the space of linearly correlated fuzzy numbers*, *Fuzzy Sets and Systems*, (2020).

- [23] L. STEFANINI, *A generalization of hukuhara difference and division for interval and fuzzy arithmetic*, Fuzzy sets and systems, 161 (2010), pp. 1564–1584.
- [24] E. SZMIDT AND J. KACPRZYK, *Distances between intuitionistic fuzzy sets*, Fuzzy Sets and Systems, 114 (2000), pp. 505–518.
- [25] C.-H. WANG, W.-Y. WANG, T.-T. LEE, AND P.-S. TSENG, *Fuzzy b-spline membership function (bmf) and its applications in fuzzy-neural control*, IEEE transactions on systems, man, and cybernetics, 25 (1995), pp. 841–851.
- [26] H. WANG AND C. ZHENG, *A class of fuzzy numbers induced by probability density functions and their arithmetic operations*, Fuzzy Sets and Systems, 467 (2023), p. 108581.
- [27] L. X. WANG, *Course In Fuzzy Systems and Control*, A, Prentice-Hall, Inc., 1996.
- [28] B. WEN, W. GU, B. YANG, H. LI, AND X. CHEN, *A novel approach for fnlp with piecewise linear membership functions*, Chemometrics and Intelligent Laboratory Systems, 191 (2019), pp. 88–95.
- [29] H.-C. WU, *Decomposition and construction of fuzzy sets and their applications to the arithmetic operations on fuzzy quantities*, Fuzzy Sets and Systems, 233 (2013), pp. 1–25.
- [30] ———, *Arithmetic operations of non-normal fuzzy sets using gradual numbers*, Fuzzy Sets and Systems, 399 (2020), pp. 1–19.
- [31] Z. XU, *Intuitionistic fuzzy aggregation operators*, IEEE Transactions on fuzzy systems, 15 (2007), pp. 1179–1187.
- [32] L. ZADEH, *The concept of a linguistic variable and its application to approximate reasoning—i*, Information Sciences, 8 (1975), pp. 199–249.
- [33] L. A. ZADEH, *Fuzzy sets, information and control*, vol, 8 (1965), pp. 338–353.